

Centre de Referència en Economia Analítica

Barcelona Economics Working Paper Series

Working Paper nº 269

Regions of Rationality: Maps for bounded agents

Robin M. Hogarth and Natalia Karelaia

October, 2005

Regions of rationality: Maps for bounded agents

Robin M. Hogarth^{a*} & Natalia Karelaia^b

ICREA^a & Universitat Pompeu Fabra^a, Barcelona

H. E. C., Lausanne,^b Switzerland

October 2005

* The authors are grateful for feedback received at workshops at the Center for Decision Research at the University of Chicago, Universitat Pompeu Fabra, Carnegie-Mellon University, the Haas School of Business at the University of California, Berkeley, and Toulouse University. This research was financed partially by a grant from the Spanish Ministerio de Educación y Ciencia.

For correspondence, please contact Robin M. Hogarth at Universitat Pompeu Fabra, Department of Economics and Business, Ramon Trias Fargas 25-27, 08005, Barcelona, Spain. (Tel: +34 93 542 2561, Fax + 34 93 542 1746). Email: robin.hogarth@upf.edu, natalia.karelaia@unil.ch

Abstract

An important problem in descriptive and prescriptive research in decision making is to identify “regions of rationality,” i.e., the areas for which simple, heuristic models are and are not effective. To map the contours of such regions, we derive probabilities that models identify the best of m alternatives ($m \geq 2$) characterized by k attributes ($k \geq 1$). The models include a single variable (lexicographic), variations of elimination-by-aspects, equal weighting, hybrids of the preceding, and models exploiting dominance. We compare all with multiple regression. We illustrate the theory with twenty simulated and four empirical datasets. Fits between predictions and realizations are excellent. However, the terrain mapped by our work is complex and no single model is “best.” We further provide an overview by regressing the performance of the different models on factors characterizing environments. We conclude by outlining how our work can be extended to exploring the effects of different loss functions as well as suggesting further topics for future research.

Keywords: Decision making, Bounded rationality, Lexicographic rules, Choice theory.

JEL classification: D81, M10.

In his autobiography, Herbert Simon (1991) used the metaphor of a maze to characterize a person's life. In this metaphor, people are continually faced by choices involving two or more alternatives, the outcomes of which cannot be perfectly predicted from the information available prior to choosing.¹ Extending this metaphor, the maze of choices a person faces can be thought of as a journey that crosses different regions varying in the types of questions posed.

If endowed with unbounded rationality, one could simply calculate the optimal responses for all decisions. However, following Simon's insights, the bounded nature of human cognitive capacities necessarily leads to following *satisficing* mechanisms. Fortunately, satisficing does not imply unsatisfactory outcomes if the type of response used is appropriate to the region in which choice is exercised. But it also raises the issue of facing the consequences of inappropriate choices.

In this paper, we characterize the maze of choices that people face as involving different "regions of rationality" where success depends on identifying decision rules that are appropriate to each region. In some regions, for example, the simplest random choice rule might be sufficient (e.g., when choosing a lottery ticket). In other regions, returns to computationally demanding algorithms are potentially important (e.g., planning production in an oil refinery). What people need therefore is knowledge – or maps – that indicate the demand for rationality in different regions. In particular, since attention is the scarce resource (Simon, 1978), it is critical to know what and how much information should be sought to make decisions in different regions.

The purpose of this paper is to contribute to defining maps that characterize regions of rationality for common decisions problems. This topic is important for both descriptive and prescriptive reasons. For the former, there is a great need to

¹ As Simon (1991) points out, this metaphor also underlies his classic (1956) paper on what an organism needs to be able to choose effectively in given environments.

understand the conditions under which simple, boundedly rational decision rules are and are not effective (see below). At the same time, this knowledge is critical for prescribing when people should use such rules, i.e., as decision aids. Specifically, we consider decisions between two or more alternatives based on information that is probabilistically related to the criterion of choice. The structure of these tasks can be conceptualized as involving either multiple-cue prediction or multi-attribute choice and, as such, is common. In all cases, we construct theoretical models that predict the effectiveness in different regions of several, simple choice rules – or heuristics (see below) – thereby mapping the contours in which the various models are more or less successful.

Following Simon's initial insights, the interest in describing the implications of simple models of decision making has grown exponentially over the last five decades (see, e.g., Conlisk, 1996; Goldstein & Hogarth, 1997; Kahneman, 2003; Koehler & Harvey, 2004). An important source of controversy in this research has centered on the extent to which the simple rules or "heuristics" that people use for making decisions are effective. In particular, great interest was stimulated by research on so-called "heuristics and biases" (Kahneman, Slovic, & Tversky, 1982) that demonstrated how simple (or less than fully rational) processes produce outcomes that deviate from normative prescriptions. Similarly, much work demonstrated that simple, statistical decision rules have superior predictive performance relative to unaided human judgment in a wide range of tasks (see, e.g., Dawes, Faust, & Meehl, 1989; Kleinmuntz, 1990).

An alternative view is that people possess a repertoire of boundedly rational decision rules that they apply in specific circumstances (Gigerenzer & Selten, 2001). Thus, heuristics can also produce appropriate responses. Specifically, Gigerenzer and

his colleagues have demonstrated how what they call “fast and frugal rules” can rival the predictive ability of complex algorithms (Gigerenzer, Todd, & the ABC Research Group, 1999). In their terms, bounded rationality can produce “ecologically rational” behavior, i.e., behavior that is appropriate in its “niche” but does not assume an underlying optimization model. What is unclear from this work, however, is where these niches are located in the regions of rationality.

Reviewing the empirical evidence, it is clear that there are occasions when heuristic rules violate normative prescriptions as well as occasions when simple rules lead to surprisingly successful outcomes. The role of theory, therefore, is to specify the circumstances in which both kinds of results occur or, to use the metaphor of this paper, to map the regions of rationality.

Our goal is to illuminate this issue and our approach is theoretical. It involves specifying analytical models for simple processes that can be used for either multi-attribute choice or multiple-cue prediction. Specifically, we derive probabilities that these models will correctly select the best of m alternatives ($m \geq 2$) based on k attributes or cues ($k \geq 1$). We also compare their effectiveness with optimizing and naïve benchmarks. The theoretical development enables the assessment of important environmental factors such as differential cue validities, inter-correlations of attributes, whether attributes/cues are measured by continuous or binary variables, levels of error in data, *and* the interactions between these factors.

This paper is organized as follows. In section I, we briefly review relevant literature. Next, in section II we specify the models we examine. In section III, we consider models based on continuous variables and derive conditions for choosing the best of three alternatives using a single variable prior to generalizing the number of alternatives and the use of different models. In section IV, we derive analogous

conditions for models based on binary attributes or cues. In section V, we test our theories on twenty simulated and four empirical datasets and find excellent fits between predictions and realizations. We also provide an overview by regressing the performance of the different models on factors characterizing environments. Our results emphasize that relative model performance is a complex function of several factors and that theoretical models are needed to understand this complexity, i.e., to map the regions of rationality. For example, we identify regions where different models do and do not exhibit similar performance. At the same time, our results are consistent with some general trends that have been demonstrated previously in simulations (e.g., effects of inter-correlation among predictor variables). We also identify new regions where “less is more” (i.e., predictions are improved if less information is used). Finally, in section VI we provide concluding comments as well as suggestions for further research.

I. Evidence on the predictive effectiveness of simple models

Interest in the efficacy of simple models for decision making has existed for some time with, in particular, numerous empirical demonstrations of how models based on simple equal (or unit) weighting schemes predict as well as or more accurately than more complex algorithms such as multiple regression (see, e.g., Dawes & Corrigan, 1974; Dawes, 1979). Gigerenzer and Goldstein (1996) have further shown how a simple, non-compensatory lexicographic model that uses binary cues (“take the best” or TTB) is surprisingly accurate in predicting the better of two alternatives across several empirical datasets and outperforms the compensatory, equal weighting (EW) model (Gigerenzer et al., 1999).

Other studies have used simulation. Payne, Bettman and Johnson (1993), for example, explored tradeoffs between effort and accuracy. Using continuous variables and a weighted additive model as the criterion, they demonstrated the effects on simple model performance of two important environmental variables, dispersion in the weighting of variables and the extent to which choices involved dominance. (See also Thorngate, 1980). Based on conceptual considerations, Shanteau and Thomas (2000) defined environments as “friendly” or “unfriendly” to different models and also demonstrated these effects through simulations.

More recently, Fasolo, McClelland, and Todd (in press) examined multi-attribute choice in a simulation using continuous variables (involving 21 options characterized by six attributes). Their goal was to assess how well choices by models with differing numbers of attributes could match total utility and, in doing so, they varied levels of average inter-correlations among the attributes and types of weighting functions. Results showed important effects for both. With differential weighting, one attribute was sufficient to capture at least 90% of total utility. With positive inter-correlation among attributes, there was little difference between equal and differential weighting. With negative inter-correlation, however, equal weighting was sensitive to the number of attributes used (the more, the better).

Despite these empirical demonstrations involving simulated and real data, research to date has generally lacked theoretical models for understanding how characteristics of models interact with those of environments. Some work has, however, considered specific cases. Einhorn and Hogarth (1975), for example, provided a theoretical rationale for the effectiveness of equal weighting relative to multiple regression. Martignon and Hoffrage (1999; 2002) and Katsikopoulos and Martignon (2003) explored the conditions under which TTB or equal weighting

should be preferred in binary choice. Hogarth and Karelaia (2004; in press) and Baucells, Carrasco, and Hogarth (2005) have examined why TTB and other simple models perform well with binary attributes in error-free environments. And, Hogarth and Karelaia (2005) provided a theoretical analysis for the special case of binary choice with continuous attributes.

II. Models considered

Whereas the essence of fully rational models involves choosing or predicting by optimally combining all relevant evidence, heuristic models are characterized by the use of limited subsets of the same information and/or simplifying combination rules (e.g., equal weighting of variables). The heuristic models we examine (see Table 1) reflect these considerations and can be classified into three categories: (A) models based on single variables or subsets of the available information; (B) equal weighting models; and (C) hybrid models that combine characteristics of the two preceding categories. In addition, we consider lower and upper benchmark models: (D) simple models that exploit dominance (see comments below); and (E) multiple regression (see also comments below).

We further examine how the type of data affects model performance by including, where possible, versions of the models based on both continuous and binary attributes/cues.² Generally speaking, we would expect models based on continuous variables to outperform their binary counterparts. However, what is not clear *a priori* is the size of such differences and how these might vary under different

² In our simulations and empirical work, we generate binary variables by median splits of the continuous variables and in this manner make direct comparisons between results based on binary and continuous variables.

conditions. We indicate the use of the two kinds of data for the same models by suffixes: $-c$ for “continuous,” and $-b$ for “binary”.

Since most of the models we consider have been considered in the literature (see previous section), we limit discussion here to making a few links. First, the DEBA model (number 3) is a deterministic version of Tversky’s (1972) elimination-by-aspects (EBA) model. For binary choice, this model is identical to the TTB model of Gigerenzer and Goldstein (1996). Variables used as attributes/cues for this model are binary in nature and, although the amount of information consulted by this model for each choice varies according to the characteristics of the alternatives, many decisions are based on a single attribute. In the continuous case, this is best matched by the single variable model (SV, number 1) which is equivalent to the lexicographic model investigated by Payne et al. (1993).

Second, with binary variables as cues/attributes, the EW model predicts frequent ties between alternatives. However, rather than resolving such choices at random, we use hybrid models that exploit partial knowledge. Specifically, EW/DEBA and EW/SVb are models that, first, attempt to choose according to EW. If this results in a tie, DEBA or SVb is used as a tie-breaker (see also Hogarth & Karelaia, in press).

Third, it is illuminating to compare the performance of simple heuristics with benchmarks. For lower or “naïve” benchmarks, we include two models that simply exploit dominance, Domran (DR), numbers 8 and 9. (Simply stated, choose an alternative if it dominates the other(s). If not, choose at random.) As an upper or “normative/sophisticated” benchmark, we use multiple regression (models 10 and 11).³

³ We are fully aware that multiple regression is not necessarily “the” optimal model for all tasks.

It is important to emphasize that the models differ in the demands they make on cognitive resources, specifically on prior knowledge and the amount of information to be processed. We therefore indicate, on the right of Table 1, differential requirements in terms of prior information, information to consult, calculations, and numbers of comparisons to be made (minimum to maximum). For example, Table 1 shows that the EW and DR models require no prior information other than the signs of the zero-order correlations between the cues and the criterion (this is a minimum requirement). On the other hand, the lexicographic, DEBA, and hybrid models need to know which cue(s) is(are) most important. Against this, the lexicographic and DEBA models do not necessarily use all cues and require no calculations. The cost of DR models lies mainly in the number of comparisons that have to be made.

Insert Table 1 about here

In this paper, we concentrate on accuracy or the probabilities that models make appropriate choices/predictions. However, and as demonstrated by Payne et al. (1993), it is important to bear in mind that heuristic models differ in their information processing costs.

Our goal is to develop theoretical models that predict model performance across different environments. However, based on the characteristics of the models, two hypotheses can be suggested. First (as noted above), we would expect models based on continuous variables to outperform their binary counterparts. Second, models that resolve ties of other models would be expected to be more accurate than the latter. Hence DEBA should be more accurate than SVb, and EW/DEBA and

EW/SVb more accurate than EWb. However, whether DEBA is more accurate than SVc will depend on environmental characteristics.

A priori, three types of environmental variables can be expected to affect absolute and relative model performance. These are, first, the distribution of “true” cue validities⁴ (i.e., how the environment weights different variables, cf., Payne et al., 1993); second, the level of redundancy or inter-correlation among the cues; and third, the level of “noise” in the environment (i.e., its inherent predictability). Of these factors, increasing noise will undoubtedly decrease performance of all models and, by extension, differences between the models. However, apart from this main effect, it is difficult to intuit how all other factors will combine to determine absolute and relative model performances. To achieve this, we need to develop appropriate theory for each of our models.

III. Models with continuous variables

Choosing the best using a single variable (SV). For expository reasons, we consider first the case of selecting the best of three alternatives using a single variable (SV). Specifically, imagine choosing from a distribution characterized by two correlated random variables, one of which is a criterion, Y , and the other an attribute, X . Furthermore, assume that alternative A is preferred over alternatives B and C if $y_a > y_b$ and $y_a > y_c$.⁵ Now, imagine that the only information about A, B, and C are the values that they exhibit on the attribute, X . Denote these specific values by x_a , x_b , and x_c , respectively. Without loss of generality, assume that $x_a > x_b$ and $x_a > x_c$ and that the decision rule is to choose the alternative with the largest value of X , i.e., in

⁴ The cue validity for a particular cue/attribute is defined by its correlation with the criterion.

⁵ We denote random variables by upper case letters, e.g., Y and X , and specific values or realizations by lower case letters, e.g., y and x . As an exception, we use lower case Greek letters to denote random error variables, e.g., ϵ .

this case A. The probability that A is in fact the correct choice can therefore be characterized by the joint probability that $Y_a > Y_b$ given that $x_a > x_b$ and $Y_a > Y_c$ conditioned on $x_a > x_c$, in other words, $P\{(Y_a > Y_b | X_a = x_a > X_b = x_b) \cap (Y_a > Y_c | X_a = x_a > X_c = x_c)\}$.

To determine this probability, assume that Y and X are both standardized normal variables, i.e., both are $N(0,1)$. Moreover, the two variables are positively correlated (if they are negatively correlated, simply multiply one by -1). Denote the correlation by the parameter ρ_{yx} , ($\rho_{yx} > 0$). Given these facts, it is possible to represent Y_a , Y_b , and Y_c by the equations:

$$Y_a = \rho_{yx}X_a + \varepsilon_a \quad (1)$$

$$Y_b = \rho_{yx}X_b + \varepsilon_b \quad (2)$$

$$\text{and } Y_c = \rho_{yx}X_c + \varepsilon_c \quad (3)$$

where ε_a , ε_b and, ε_c are normally distributed error terms, each with mean of 0 and variance of $(1 - \rho_{yx}^2)$, independent of each other and of X_a , X_b , and X_c .

Using equations (1), (2), and (3) the differences between Y_a and Y_b , on the one hand, and Y_a and Y_c , on the other, can be written as

$$Y_a - Y_b = \rho_{yx}(X_a - X_b) + (\varepsilon_a - \varepsilon_b) \quad (4)$$

and

$$Y_a - Y_c = \rho_{yx}(X_a - X_c) + (\varepsilon_a - \varepsilon_c) \quad (5)$$

Thus, $Y_a > Y_b$ and $Y_a > Y_c$ if

$$\rho_{yx}(X_a - X_b) > \varepsilon_b - \varepsilon_a \quad (6)$$

and

$$\rho_{yx}(X_a - X_c) > \varepsilon_c - \varepsilon_a \quad (7)$$

$P\{(Y_a > Y_b | X_a = x_a > X_b = x_b) \cap (Y_a > Y_c | X_a = x_a > X_c = x_c)\}$ can now be reframed as the probability that *both* the right hand side of (6) is smaller than $\rho_{yx}(X_a - X_b)$ and the right hand side of (7) is smaller than $\rho_{yx}(X_a - X_c)$. As can be seen, these latter terms are the products of ρ_{yx} , the correlation between Y and X , and the differences between X_a and X_b , and X_a and X_c . In other words, the larger the correlation between Y and X , and the larger the differences between X_a and X_b , and X_a and X_c , the greater $P\{(Y_a > Y_b | X_a = x_a > X_b = x_b) \cap (Y_a > Y_c | X_a = x_a > X_c = x_c)\} =$

$$P\{(\varepsilon_b - \varepsilon_a < \rho_{yx}(x_a - x_b)) \cap (\varepsilon_c - \varepsilon_a < \rho_{yx}(x_a - x_c))\} \quad (8)$$

To determine this probability, we make use of the facts that the differences between the error terms, $(\varepsilon_b - \varepsilon_a)$ and $(\varepsilon_c - \varepsilon_a)$, are both normally distributed with means of 0 and variances of $2(1 - \rho_{yx}^2)$. Standardizing $(\varepsilon_b - \varepsilon_a)$ and $(\varepsilon_c - \varepsilon_a)$, we can re-express equation (8) as

$$P\{(\varepsilon_b - \varepsilon_a < \rho_{yx}(x_a - x_b)) \cap (\varepsilon_c - \varepsilon_a < \rho_{yx}(x_a - x_c))\} = P\left\{\left(z_1 < \frac{\rho_{yx}(x_a - x_b)}{\sqrt{2(1 - \rho_{yx}^2)}}\right) \cap \left(z_2 < \frac{\rho_{yx}(x_a - x_c)}{\sqrt{2(1 - \rho_{yx}^2)}}\right)\right\} \quad (9)$$

where z_1 and z_2 are standardized normal variables with means of 0 and variances of 1. Moreover, z_1 and z_2 jointly follow a bivariate normal distribution. Therefore, the target probability (9) can be written as

$$\int_{-\infty}^{l_{ab}} \int_{-\infty}^{l_{ac}} \frac{1}{2\pi\sigma_{z_1}\sigma_{z_2}\sqrt{1-\rho^2}} e^{-\frac{z}{2(1-\rho^2)}} dz_1 dz_2 \quad (10)$$

where $z = \frac{z_1^2}{\sigma_{z_1}^2} - \frac{2\rho z_1 z_2}{\sigma_{z_1}\sigma_{z_2}} + \frac{z_2^2}{\sigma_{z_2}^2}$; $l_{ab} = \frac{\rho_{yx}(x_a - x_b)}{\sqrt{2(1 - \rho_{yx}^2)}}$; $l_{ac} = \frac{\rho_{yx}(x_a - x_c)}{\sqrt{2(1 - \rho_{yx}^2)}}$;

$\sigma_{z_1} = \sigma_{z_2} = 1$; and $\rho = \sigma_{z_1, z_2}$.

In Appendix A, we show that $\rho = 1/2$. Thus, we can write

$$P\{(Y_a > Y_b | X_a = x_a > X_b = x_b) \cap (Y_a > Y_c | X_a = x_a > X_c = x_c)\} = \int_{-\infty}^{l_{ab}} \int_{-\infty}^{l_{ac}} \frac{1}{\pi\sqrt{3}} e^{-\frac{2}{3}(z_1^2 - z_1 z_2 + z_2^2)} dz_1 dz_2 \quad (11)$$

Figure 1 illustrates the probabilities of SV correctly choosing the best of three alternatives for different values of $(X_a - X_b)$ and $(X_a - X_c)$. In the panel on the left, (a), $(x_a - x_c)$ is held constant at a low value of 0.3; on the right, (b), it is held constant at a high value, 2.0. The lines in the figures reflect the effects of combining these fixed levels with different values of $(x_a - x_b)$, from 0.1 to 3.0. As can be observed, if one of the two differences, $(x_a - x_b)$ or $(x_a - x_c)$, is small, the probability of a correct choice varies between 0.4 and 0.5. However, as both grow larger, so do the corresponding probabilities.

 Insert Figure 1 about here

To generalize the above, assume that there are m ($m > 3$) alternatives from which to choose and that each has a specific X value, x_l , $l = 1, \dots, m$. Without loss of generality, assume that x_l has the largest value and we wish to know the probability that the corresponding alternative has the largest value on the criterion. Generalizing from the above, this probability can be calculated using properties of the multivariate normal distribution and, in this case, can be written,

$$\int_{-\infty}^{d_1^*} \dots \int_{-\infty}^{d_{m-1}^*} \varphi(z | \mu_z, V_z) dz_1 \dots dz_{m-1} = \int_{-\infty}^{d_1^*} \dots \int_{-\infty}^{d_{m-1}^*} \frac{|V_z|^{1/2}}{(2\pi)^{(m-1)/2}} e^{-\frac{1}{2}z'V_z^{-1}z} dz_1 \dots dz_{m-1} \quad (12)$$

where $d_i^* = \frac{\rho_{yx} d_i}{\sqrt{2(1-\rho_{yx}^2)}}$ for $i = \overline{1, m-1}$, the elements of $z' = (z_1, z_2, \dots, z_{m-1})$ are jointly

distributed normal variables, with means of zero and variances of one, and V_z^{-1} is the inverse of the $(m-1) \times (m-1)$ variance-covariance matrix where each diagonal element is equal to 1 and all off-diagonal elements equal $\frac{1}{2}$ (see Appendix A). In Appendix B we derive the analytical expression for the probability of selecting the optimal choice among four alternatives by using just one variable. For binary choice, that is, when $m = 2$, analogous derivations lead to similar expressions to those shown above (see Hogarth & Karelaia, 2005).

Overall probabilities. The probabilities given above are those associated with particular observations, i.e., that A is larger than B and C given that a specific value, x_a , exceeds specific values x_b and x_c . However, it is also instructive to consider the overall expected accuracy of SV, i.e., the overall probability that SV makes the correct choice when sampling at random from the population of alternatives.

Overall, SV can make successful choices in three ways: selecting A when x_a is bigger than x_b and x_c ; selecting B when x_b is bigger than x_a and x_c ; and selecting C when x_c is bigger than x_a and x_b . The probabilities of these events are,

$$P\{((X_a > X_b) \cap (X_a > X_c)) \cap ((Y_a > Y_b) \cap (Y_a > Y_c))\},$$

$$P\{((X_b > X_a) \cap (X_b > X_c)) \cap ((Y_b > Y_a) \cap (Y_b > Y_c))\}, \text{ and}$$

$$P\{((X_c > X_a) \cap (X_c > X_b)) \cap ((Y_c > Y_a) \cap (Y_c > Y_b))\}$$

respectively, and the overall probability is the sum of the three terms. However, since each of the terms is equal to the others, the sum can be re-expressed as $3P\{((X_a > X_b) \cap (X_a > X_c)) \cap ((Y_a > Y_b) \cap (Y_a > Y_c))\}$.

To derive analytically the overall probability of correct choice by SV when sampling at random from the underlying population of alternatives, the latter expression should be integrated across all possible values that can be taken by $D_1 = X_a - X_b > 0$, and $D_2 = X_a - X_c > 0$. That is

$$3 \int_0^{\infty} \int_0^{\infty} \varphi(d|\mu_d, V_d) \left[\int_{-\infty}^{d_1^*} \int_{-\infty}^{d_2^*} \varphi(z|\mu_z, V_z) dz_1 dz_2 \right] dd_1 dd_2 \quad (13)$$

where $z' = (z_1, z_2)$, $V_z = \begin{pmatrix} 1 & 1/2 \\ 1/2 & 1 \end{pmatrix}$, $d' = (d_1, d_2)$, $V_d = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$, $d_1^* = \frac{\rho_{yx} d_1}{\sqrt{2(1-\rho_{yx}^2)}}$,

and $d_2^* = \frac{\rho_{yx} d_2}{\sqrt{2(1-\rho_{yx}^2)}}$. (In Table 3, discussed below, we generalize these formulas for

choosing one of m alternatives.)

Equal weighting (EW) and multiple regression (MR). What are the predictive accuracies of models that make use of several, k , cues or variables, $k > 1$? We consider two models that have often been used in the literature. One is equal weighting (EW – see Dawes & Corrigan, 1974; Einhorn & Hogarth, 1975). The other is multiple regression (MR). To analyze these models, assume that the criterion variable, Y , can be expressed as a function

$$Y = f(X_1, X_2, \dots, X_k) \quad (14)$$

where the k predictor variables are multivariate normal, each with mean of 0 and standard deviation of 1. For EW, the predicted Y value associated with any vector of

observed x 's is equal to $\frac{1}{k} \sum_{j=1}^k x_j$ or \bar{x} . Similarly, the analogous prediction in MR is

given by $\sum_{j=1}^k b_j x_j$ or \hat{y} where the b_j 's are estimated regression coefficients. In using

these models, therefore, the decision rules are to choose according to the largest \bar{x} for EW and the largest \hat{y} value for MR.

How likely are EW and MR to make the correct choice? Following the same rationale as the single variable (SV) case, we show in Table 2 the formulas used in deriving the analogous probabilities for EW and MR (as well as SV) when choosing the best of three alternatives using the properties of the bivariate normal distribution. These are the initial equations (corresponding to equations 1, 2, and 3), the accuracy conditions (corresponding to equations 6 and 7), the relevant error variances, and finally the upper limits of integration, i.e. l_{ab} and l_{ac} , used to calculate probabilities when applying equation (11).

 Insert Tables 2 & 3 about here

Similarly, when calculating the probabilities of choosing correctly between four or more alternatives for EW and MR, we can apply the multivariate normal distribution in analogous fashion to that of SV (cf. equation 12 above and Appendix B).

In Table 3, we present the formulas for the overall expected accuracy of EW and MR in a given environment or population, analogous to those for SV, i.e. to equation (13), for choosing one of m alternatives. In particular, we present the elements that are specific for different models, such as the variance-covariance matrix, V_d , and the upper integration limits of integration, d_i^* .

IV. Models with binary variables

To discuss expected predictive performance of models based on binary variables, we first assume that the dependent variable, Y , can be thought of as being generated by a linear model of the form

$$Y = a + \sum_{j=1}^k \gamma_j W_j + \zeta \quad (15)$$

where $W_j = 0, 1$ are the binary variables ($j = 1, \dots, k$), the γ_j are weighting parameters and ζ is a normally distributed error term (see also below).

To derive theoretical predictions for models using binary variables, we adopt a similar approach to that used with continuous variables. We therefore focus on issues that differ between the continuous and binary cases.

Choosing the best using a single binary variable (SVb). Assuming that $w_a > w_b$ and $w_a > w_c$, the probability that SVb chooses correctly between three alternatives, A, B, and C is $P\{(Y_a > Y_b | W_a = w_a > W_b = w_b) \cap (Y_a > Y_c | W_a = w_a > W_c = w_c)\}$.

To determine this probability, recall that Y is a standardized normal variable $N(0,1)$. The binary variable, W , however, only takes values of 0 and 1 and thus has a mean of 0.5 and standard deviation, σ_w , of 0.5.⁶ Denoting the correlation between Y and W by ρ_{yw} , ($\rho_{yw} > 0$), we can express Y by

$$Y = a_{SVb} + \frac{\rho_{yw}}{\sigma_w} W + \zeta$$

or, simply,
$$Y = a_{SVb} + 2\rho_{yw}W + \zeta \quad (16)$$

where ζ is a normally distributed error term $N(0, 1 - \rho_{yw}^2)$.⁷

Proceeding in similar fashion to the continuous case, we obtain the expression for the probability of SVb predicting correctly:

$$\int_{-\infty}^{h_{ab}} \int_{-\infty}^{h_{ac}} \frac{1}{\pi\sqrt{3}} e^{-\frac{2}{3}(z_1^2 - z_1 z_2 + z_2^2)} dz_1 dz_2 \quad (17)$$

⁶ Recall that binary variables are created by median splits of continuous variables.

⁷ Since $E(Y) = 0$, it follows that the intercept $a_{SVb} = -\rho_{yw}$.

where $h_{ab} = \frac{2\rho_{yw}(w_a - w_b)}{\sqrt{2(1 - \rho_{yw}^2)}} ; h_{ac} = \frac{2\rho_{yw}(w_a - w_c)}{\sqrt{2(1 - \rho_{yw}^2)}} .$

Since both $(w_a - w_b)$ and $(w_a - w_c)$ are equal to one, the two upper integration limits

are the same: $h_{ab} = h_{ac} = \frac{2\rho_{yw}}{\sqrt{2(1 - \rho_{yw}^2)}} .$

As can be seen, the only difference between the theoretical expressions for the continuous and binary cases lies in the formulas for the upper limits of integration. Therefore, generalizing the above for choices among m ($m > 3$) alternatives is analogous to that for the continuous case.

Following the same rationale, we can derive the formulas for the probabilities for EWb and MRb when choosing the best of three alternatives using binary variables. In Table 4, we present the initial equations for these models (corresponding to equation 16), and the upper limits of integration, i.e. h_{ab} and h_{ac} , used to calculate probabilities when applying equation (17).

 Insert Table 4 about here

Choosing the best using DEBA with binary cues. Recall that this multi-stage model works in the following way. At the first stage, alternatives with values of 0 for the most important cue are eliminated unless all alternatives exhibit 0. If only one alternative has a value of 1, it is selected and the process terminates. If, however, more than one alternative remains, the same procedure takes place with the remaining alternatives except that the second most important cue is used. The process continues in the same manner through subsequent stages, if necessary. It stops when either only one alternative remains (i.e., the chosen alternative) or, if there is more than one

alternative but no more cues, choice is determined at random among the remaining alternatives.

The probability that a given alternative was chosen correctly by DEBA is the probability that the sequence of decisions (or eliminations) made by the model at each stage is correct. Thus, since at each stage of the model decisions are made conditional on the preceding stages, the key parameters in estimating these probabilities are the partial correlations between Y and $W_j, j = 1, \dots, k$ (i.e., controlling for previous stages). For the first stage, this is ρ_{yw_1} , for the second $\rho_{yw_2.w_1}$, for the third, $\rho_{yw_3.w_1w_2}$, and so on.⁸

For example, assume that there are three alternatives A, B, and C and that A has been chosen by a process whereby C was eliminated at the first stage and B at the third stage. Starting backwards, consider the decisions the model makes at each stage. That is, the probability that DEBA correctly selected A over B at the third stage, controlling for the elimination of C at the first stage, is $P\{(Y_a > Y_b | W_{a3} = w_{a3} > W_{b3} = w_{b3}) \cap (Y_b > Y_c | W_{b1} = w_{b1} > W_{c1} = w_{c1})\}$. This probability can be calculated by making use of the appropriate partial correlations – in this case, $\rho_{yw_3.w_1w_2}$ and ρ_{yw_1} – and adapting the single variable equations (e.g., the general equation 10⁹). At the second stage, the model makes no decision. At the first stage, it eliminates C so we need to calculate additionally the probability that A could have been correctly selected only with information available at this stage: $P\{(Y_a > Y_c | W_{a1} = w_{a1} > W_{c1} = w_{c1}) \cap (Y_c > Y_b | W_{b1} = w_{b1} > W_{c1} = w_{c1})\}$. This can be also found through an adapted expression (10), using ρ_{yw_1} . Importantly, the events

⁸ For example, $\rho_{yw_2.w_1} = \frac{\rho_{yw_2} - \rho_{yw_1}\rho_{w_1w_2}}{\sqrt{(1 - \rho_{yw_1}^2)(1 - \rho_{w_1w_2}^2)}}$.

⁹ The terms that need to be adapted in the expression (10) are the upper limits of integration and σ_{z_1, z_2} .

represented by the probability expressions for the first and third stages are disjunctive. Therefore, the probability that DEBA makes the correct decision in this case is equal to the sum of the two expressions.

Consider another example involving three alternatives A, B, and C. Assume that DEBA eliminates C at the first stage and at the third stage picks either A or B at random (this will happen if A and B are identical). Thus, the 0.5 probability that DEBA makes the correct decision at the third stage should be “discounted” by the probability that C, eliminated at the first stage, is not better than A and B. That is

$$0.5(1 - P\{(Y_c > Y_a | W_{c1} = w_{c1} > W_{a1} = w_{a1}) \cap (Y_c > Y_b | W_{c1} = w_{c1} > W_{b1} = w_{b1})\}).$$

More generally, the probability of DEBA making the correct choice has to be calculated on a case-by-case basis taking into account, at each stage, the probability that the selected alternative should be chosen over the alternative(s) eliminated at that stage using the partial correlation of the cue appropriate to the stage. Moreover, the probability for each case includes the probabilities of successful decisions at each stage. If at the final stage, there are two or more alternatives, the appropriate random probability is adjusted by the probability that correct decisions were taken at previous stages (see, e.g., the example above).¹⁰

Choosing the best using the EW-SV model with binary cues (EW-SVb). The first stage of this model uses EWb. If a single alternative is chosen, the probability of it being correct is found by applying the formula for EWb. If two or more alternatives are tied, a second stage consists of selecting the alternative favored by the first cue. To calculate the probability that this is correct, one needs to calculate the joint probability that the selected alternative (a) is larger than the alternatives eliminated at

¹⁰ In this section, we have only indicated the general strategy for calculating relevant probabilities for DEBA. The details involve repeated applications of the same probability theory principles applied in many different situations (Karelaia & Hogarth, in preparation).

the first stage, and (b) larger than the other alternatives considered at the second stage. (To calculate these probabilities, use is made of the appropriate analogs to equation 17). Any ties remaining after stage two are resolved at random with a corresponding adjustment being made to the probability calculations.

Choosing the best using the EW-DEBA model with binary cues (EW-DEBA).

This model starts as EWb. If EWb chooses one alternative, the probability of correct choice of the model coincides with that for EWb. If two or more alternatives are tied, the DEBA model is used to choose between the remaining alternatives and probabilities are calculated accordingly (see above).

V. Empirical evidence

Our equations provide exact theoretical probabilities for assessing performance of the different models in specified conditions, i.e., to map the contours of the regions of rationality. However, several factors affect absolute and relative performance levels of the models (e.g., cue validities, inter-correlation among variables, continuous vs. binary variables, error), and it is difficult to assess their importance simply by inspecting the formulas.

We therefore use both simulated and empirical data to illuminate model performance under different conditions. Real data have the advantage of testing the theory in specific, albeit limited environments. Simulated data, on the other hand, facilitate testing model predictions over a wide range of environments. We first consider the simulated data.

Simulation design and method. The simulation design used for choosing the best from two, three and four alternatives is presented in Table 5. Overall, we specified 20 different populations that are subdivided into four sets or cases – A, B, C,

and D – each of which contains five sub-cases (labeled 1, 2, 3, 4, and 5). Since, a priori, several factors might be thought important, we would have liked to vary these orthogonally. However, correlations between factors restrict implementing a fully systematic design. We therefore varied some factors at the level of the cases (A, B, C, and D) and others across sub-cases (i.e., within A, B, C, and D).

At the level of cases, A and B involved three cues or attributes whereas cases C and D involved five. Cases A and C had little or no inter-cue correlation; cases B and D had moderate to high intercorrelation.

Insert Table 5 about here

Across sub-cases (i.e., from 1 through 5 within each of A, B, C, and D), we varied: (1) the variability of cue validities (maximum less minimum); (2) the validity of the first (i.e., most important) cue; (3) average validity; and (4) the correlation between y and \bar{x} . For all, values increase from the sub-cases 1 through 5. As a consequence, the R^2 on initial fit for MR also increases across sub-cases. This implies that the sub-cases 1 involve high levels of error whereas the sub-cases 5 are, in principle, quite predictable environments. Sub-cases 2, 3, and 4 fall between these extremes.

To conduct the simulation, we defined 20 sets of standardized multivariate normal distributions with the parameters specified in Table 5 and generated samples of size 40 from each of these populations. The observations in each sample were split at random on a 50/50 basis into fitting and prediction sub-samples and model parameters were estimated on the fitting sub-sample. Two, three or four alternatives (as appropriate) were then drawn at random from this sub-sample and, using the estimated model parameters, probabilities of correctly selecting the best of these

specific alternatives were calculated. This was then compared to what actually happened, that is, on a “fitting” basis. Next, alternatives were drawn at random from the prediction sub-sample, relevant probabilities calculated using the parameters from the fitting sub-sample, and predictions compared to realizations. This exercise was repeated 5,000 times (for each of the choices involving two, three, and four alternatives).¹¹

The above describes the procedure used for continuous data. For models using binary data, we followed exactly the same procedures except that predictor variables only took values of 0 or 1. Specifically, since we were sampling continuous normalized variables, we created binary variables by median splits (i.e., binary variables were set to 0 for negative values of continuous variables and 1 for non-negative values). Thus, if one estimates the parameters of the 20 populations for the binary data, the estimates differ systematically from their continuous counterparts shown in Table 5 (they are smaller). However, we do not display the binary parameter estimates since the parameters in Table 5 represent the process that generated the data.

Simulation results. Tables 6, 7, and 8 present the results of the simulations for the choice of best of two, three, and four alternatives, respectively. Figure 2 presents some selected outcomes from the case involving best of three (Table 7). Results reported here are limited to predictions and realizations for the holdout samples (i.e., tests of cross-validation).¹²

¹¹ A possible criticism of our predictive tests of the single variable models (SVc, SVb, and DEBA) is that we did not use the sampling process to determine the most important variable (for SVc and SVb) nor the rank orders of the cue validities (for DEBA). Instead, we endowed the models with the appropriate knowledge. However, in subsequent simulations we have found that with sample sizes of 20 (as here) the net effect of failing to identify the most important variable is quite small. Similarly, as long as DEBA correctly identifies the most important variable, net differences are also small (Hogarth & Karelaia, in press).

¹² As might be expected by examining the success of the models on cross-validation, the fitting exercise produced almost perfect matches between samples and models.

To simplify reading the tables, note that realizations are underlined (e.g., 62), and that the largest realization for each population (i.e., per column) is presented in bold (e.g., **65**). In addition, when MRc is the largest, we also denote the second largest in bold. We further show mean realizations for each column and row in the tables. The column means thus represent the average realizations of all models within specific populations whereas the row means characterize average model performance across populations.

We first note that, with the exception of multiple regression (MR), the match between model predictions and realizations is quite close in all cases. MR makes large errors for cases C and D and particularly when the differences between maximum and minimum cue validities are smallest (sub-cases 1 and 2). Although we used adjusted R^2 in making predictions, the adjustment was insufficient in these situations.¹³

Insert Tables 6, 7, 8 and Figure 2 about here

Qualitatively, the relative effectiveness of the models is quite similar whether one looks at the results for best of two, three, or four. What changes, of course, is the general level of performance which diminishes as the number of alternatives increases. This can be seen by comparing the columns of mean realizations of Tables 6, 7, and 8, i.e., at the extreme right hand sides of the tables.

Within each table, an initial, overall impression is the lack of large differences between the performances of the different models. However, there are systematic effects.

¹³ One can also argue with some justification that the ratio of observations to predictor variables is too small to use multiple regression (particularly for cases C and D). However, we are particularly interested in observing how well the different models work in environments where there are not many observations.

First, consider whether predictor variables are binary or continuous. The use of binary as opposed to continuous variables implies a loss of information. As such, we expected that models based on continuous variables would predict better than their binary counterparts. Indeed, this is always the case in three direct comparisons: SVc vs. SVb, EWc vs. EWb, and MRc vs. MRb. Specifically, note that SVb and EWb are both handicapped relative to SVc and EWc in that they necessarily predict many ties that are resolved at random. Thus, so long as the knowledge in SVc and EWc implies better than random predictions, models based on continuous variables are favored.

On the other hand, the performance of DRb dominates DRc for cases A, C, and D with performance being quite similar for case B (for best of two, three, and four). It would appear that DRb exploits more cases of “apparent” dominance than DRc which (as a consequence) decides more choices at random. Thus, to the extent that the “additional” dominance cases detected by DRb relative to DRc have more than a random chance of being correct, DRb outpredicts DRc. Whereas the DR models represent naïve baseline strategies, we believe this finding is important because it demonstrates how a simple strategy can exploit the structure of the environment such that more information (in the form of continuous as opposed to binary variables) does not improve performance (a so-called “less is more” effect, Goldstein & Gigerenzer, 2002; Hertwig & Todd, 2003).

Second, models that resolve ties perform better than their counterparts that are unable to do so (e.g., DEBA vs. SVb, and EW/DEBA and EW/SVb vs. EW). However, in the presence of redundancy, these differences are quite small (i.e., for cases B and D). More interesting is the comparison between SVc and DEBA. The former uses a single, continuous variable. The latter relies heavily on one binary variable but can also use others depending on circumstances. It is thus not clear

which strategy actually uses more information. However, once again, characteristics of the environment determine which strategy is more successful. SVc dominates DEBA in case B as well as for much of cases A and D. On the other hand, DEBA dominates SVc in case C.

In Figure 2, we have chosen to illustrate the performance of five models across all the environments for the choice of best of three (i.e., data from Table 7). Two of these models, DRb and MRc are depicted because they represent, respectively, naïve and “sophisticated” benchmarks. The other models, SVc, DEBA, and EWc are quite different types of heuristics (see Table 1). Both SVc and DEBA require prior knowledge of what is important (DEBA more so than SVc). However, they use little information and neither involves any computation. (DEBA, it should be recalled, also operates on binary data.) EWc, on the other hand, does not require knowledge of differential importance of variables but does use all information available and needs some computational ability.

In interpreting Figure 2, it is instructive to recall that cases A and C (on the left) represent environments with low redundancy whereas cases B and D (on the right) have higher levels of redundancy. Also within each case, the amount of noise in the environment decreases as one moves from sub-case 1 (on the left) to sub-case 5 (on the right).

As expected, in the noisier environments (sub-cases 1), the performances of all models are degraded such that differences are small. However, as error decreases (i.e., moving right toward sub-cases 5), model performances vary by environmental conditions. With low redundancy (cases A and C), there appear to be large differences in model performance. However, in the presence of redundancy (cases B and D), there are two distinct classes of models: SVc and MRc have similar

performance levels and are superior to the others. We further note that SVc is most effective in case B and also does well as environmental predictability increases in cases A and D. DEBA is never the best model but performs quite adequately in case C where EWc has the best performance. Of the benchmark models, DRb generally lags behind the other models (as would be expected). Finally, although MRc is typically one of the better models, it does not dominate in all environments.

To highlight regions of rationality, Table 9 represents the data from Figure 2 in another manner (think of Table 9 as a map!). Specifically, the performances of SVc, DEBA, EWc, and DRb are compared to the “normative” benchmark of MRc (by deducting the performance of each of the former from the latter). Thus positive (negative) entries in Table 9 indicate the amount by which the performance of MRc exceeds (falls short of) those of other models. Three cases are indicated: in the shaded areas, MRc exceeds other models by at least 5; in the unshaded areas, the standard font (e.g., 4) indicates that the MRc advantage is small (< 5) but positive; and bold, underlined font (e.g., **-2**) denotes that MRc has no advantage over the other model. The table depicts relative model performance by characteristics of regions, e.g., redundancy and noise (that is, inter-cue correlation and error, respectively).

Insert Table 9 about here

For example, recall that cases A and C involve low redundancy whereas this is not true of cases B and D. Also, for all cases, as one moves from sub-cases 1 through 5, environments involve less noise. With this in mind, one can attribute the smaller differences between models in sub-cases 1 and 2 as being mainly due to noise. Interestingly, the relative success of SVc in cases B and D (compared to A and C) seems to be the effect of inter-correlation between the other attributes, i.e.,

redundancy. Table 9 is only presented as an illustration. The data from Tables 6, 7, and 8 can clearly be used to create maps that highlight different aspects of the decision making terrain.

A further way of summarizing factors that affect the performance of the different models is to consider the regression of model performance on statistics that describe the characteristics of the 20 simulated environments (Table 5). In other words, consider regressions for each of our eleven models of the form

$$P_i = Z \underline{\delta}_i + \tau_i \quad (18)$$

where P_i is the performance realization of model i ($i = 1, \dots, 11$); Z is the $(20 \times 3) \times s$ matrix of independent variables (statistics characterizing the datasets where there are three choice situations, i.e., best of two, three, or four alternatives); $\underline{\delta}_i$ is the $s \times 1$ vector of regression coefficients; and τ_i is a normally distributed error term with constant variance, independent of Z .

To characterize the environments or datasets (the Z matrices), we chose the following variables: variability of cue validities (max less min), the validity of the most important cue (r_{yx_1}), the validity of the average of the cues ($r_{y\bar{x}}$), average inter-correlation of the cues, average validity of the cues, number of cues, and R^2 for MRc and MRb.¹⁴ We also used dummy variables to model the effects of choosing between different numbers of alternatives. Dummy1 captures the effect of choosing from three as opposed to two alternatives, and Dummy2 the additional effect of choosing from four alternatives. Results of the regression analyses are summarized in Table 10.

¹⁴ We only used R^2 as an independent variable for MRc and MRb because we thought it would be appropriate for these models. For the other models, however, it was deemed more illuminating to characterize performance by the other measures (R^2 and these other measures are correlated in different ways). It should also be noted that we used the same statistics (based on continuous variables) to characterize the environments for models using both continuous and binary variables on the grounds that the underlying environments were based on continuous variables.

Insert Table 10 about here

We used a step-wise procedure with entry (exit) thresholds for the variables of $<.05$ ($>.10$) for the probability of the F statistic. All coefficients for the models shown in Table 10 are statistically significant ($p < .001$) and all regressions fit the data well (see R^2 and estimated standard errors at the foot of Table 10). The constant term is fairly high across all models and measures the level of performance that would be expected of the models in binary choice absent information about the environment (approximately 50, i.e., from 42 to 65). Dummy1 indicates how much such performance would fall when choosing between three alternatives (between 11 and 14), and Dummy2 shows the additional drop experienced when choosing among four alternatives (between 6 and 9).

For SVc and SVb, only one other variable is significant, the correlation between the single variable and the criterion. This makes intuitive sense as does the fact that the regression coefficient is larger with continuous as opposed to binary variables (50 vs. 29). The DEBA and EW models are all heavily influenced by the correlation between the criterion and \bar{x} . Recall, however, that this correlation is itself an increasing function of average cue validity and the number of cues but decreasing in the inter-correlation between cues (see the formula in footnote 1 to Table 2). Thus, *ceteris paribus*, increasing inter-correlation between the cues reduces the absolute performance levels of these models. DEBA differs from the EW models in that the correlation of the most valid cue is a significant predictor. This matches expectations in that DEBA relies heavily on the validity of the most important cue whereas EW weights all cues equally. (We also note that the SV models weight the most valid cue more heavily than DEBA.) As to the Domran models, the interpretation of the signs of

all coefficients is not obvious. Finally, for MRc it comes as little surprise that R^2 should be so important although this variable is less salient for MRb.

A possible surprise is that variability in cue validities (maximum less minimum) was not a significant factor for most models. One might have thought, a priori, that such dispersion would have been important for DEBA (cf., Payne et al., 1993). However, this is not the case and is consistent with theoretical analyses of DEBA that show that its accuracy is relatively robust to different “weighting functions” (Hogarth & Karelaia, in press; Baucells, Carrasco, & Hogarth, 2005).

Finally, whereas the regression statistics paint an interesting picture of model performance in the particular environments observed, we caution against overgeneralization. We only observed restricted ranges of the environmental statistics (i.e., characteristics) and thus cannot comment on what might happen beyond these ranges. Our approach, however, does suggest a way to illuminate model x environment interactions.

To summarize, across all 20 environments that, inter alia, are subject to different levels of error, the relative performances of the different models were not seen to vary greatly when faced with the same tasks (e.g., choose best of three alternatives). However, there were systematic differences due to interactions between characteristics of models and environments. Thus, whereas the additional information contained in continuous as opposed to binary variables benefits some models, e.g., SV and EW, it can be detrimental to others, e.g., DR. Second, models varied in the extent to which they were affected by specific environmental characteristics. SV models, for example, depend heavily on the validity of the most valid cue whereas this only affects EW models through its impact on average cue validity. Interestingly, the validity of the average of the cues was seen to have more impact on the performance

of DEBA than the validity of the most valid cue. Average inter-correlation of predictors or redundancy tends to reduce performance of all models (except SV).

Overall, results do match some general trends noted in previous simulations (Payne et al., 1993; Fasolo et al., in press); however, patterns are not simple to describe. The value of our work, therefore, is that we now possess the means to make precise predictions for various simple, heuristic models in different environments. That is, given specified environments, we can predict *a priori* both levels of model performance and which models will be more or less effective.

An important environmental factor we did not vary was the impact of different distributions of specific kinds of alternatives. Instead, we simulated random drawings of alternatives given the population characteristics defined in Table 5. We did not, for example, skew the sampling process to include or exclude disproportionate numbers of, say, dominating or dominated alternatives (cf., Payne et al., 1993). As we have argued elsewhere (Hogarth & Karelaia, 2004), the distribution of alternatives can have important effects on both the general level of performance achieved by models as well as relative performance (some distributions, for example, are relatively “friendly” or “unfriendly” to specific models, Shanteau & Thomas, 2000). On the other hand, since our methodology can make specific predictions for each case encountered, it can easily handle the effects of sampling from different distributions of alternatives.

Empirical data. We used datasets from three different areas of activity. The first involved performance data of the 60 leading golfers in 2003 classified by the Professional Golf Association (PGA) in the USA.¹⁵ From these data (N = 60), we examined two dependent variables: “all-round ranking” and “total earnings.” The first

¹⁵ These data were obtained from the webpage <http://www.pgatour.com/stats/leaders/r/2003/120>. They are performance statistics of golfers in the main PGA Tour for 2003.

is a measure based on eight performance statistics. For our models, we chose three predictor variables that account for 67% of the variance in the criterion. These were mean numbers – across rounds played – of birdies, total scores, and putts. Since the first variable was negatively related to the criterion, it was rescaled (multiplying by -1).

Eighty-two percent of the variance in the second golf criterion, total earnings, could be explained by three variables, “number of top 10 finishes,” “all-round ranking” (the previous dependent variable), and “number of consecutive cuts.” Of these, all-round ranking was negatively correlated with the criterion and so rescaled (multiplying by -1).

The second dataset consisted of rankings of PhD economics programs in the USA on the basis of a 1993 study by the National Research Council (N = 107).¹⁶ Three variables accounted for 80% of the variance in the rankings: number of PhD’s produced by programs for the academic years 1987-88 through 1991-1992, total number of program citations in the period 1988-1992 divided by number of program faculty, and percentage of faculty with research support.

The third dataset was taken from the UK consumer organization *Which?*’s assessments of digital cameras in 2004 (N = 49).¹⁷ Three variables were found to explain 72% of the variance in total test scores: image quality, picture downloading time, and focusing.

Table 11 summarizes statistical characteristics of these datasets (analogous to the experimental design of the simulations in Table 5). As can be seen, there is little to moderate variability in cue validities (compare *Economics PhD programs* with the other datasets); all datasets have at least one highly valid cue; average inter-

¹⁶ For more details, see the webpage <http://www.phds.org>

¹⁷ For further details, see the webpage <http://sub.which.net>

correlation varies from moderate to low (*Consumer reports: Digital cameras*); and there are high correlations between the criteria and the means of the predictor variables.

The testing procedure was similar to the simulation methodology. We divided each sample at random on a 50/50 basis into fitting and testing sub-samples. Model parameters were then estimated on the fitting sub-sample and these parameters used to calculate the probabilities that the models would correctly choose the best of two, three, and four alternatives that had also been randomly drawn from the same sub-sample. This was the fitting exercise. To test the models' predictive abilities, two, three, and four alternatives were drawn at random from the second or testing sub-sample and, using the parameters estimated from the data in the first or fitting sub-sample, model probabilities were calculated for the specific cases and subsequently compared to realizations. This exercise was repeated 5,000 times such that the data reported in Table 12 shows the aggregation of all these cases. Also similar to the simulation methodology, we created binary datasets by using median splits of the continuous variables.

Insert Tables 11, 12 and Figure 3 about here

Table 12 has been constructed in similar fashion to Tables 6, 7, and 8 except that we show results for choosing the best out of two, three, and four alternatives within columns for each dataset.

Once again, there is an excellent fit between predictions and realizations.¹⁸ The largest differences occur, as before, with MR. Overall, the models have fairly high levels of performance which, naturally, diminish as the number of alternatives

¹⁸ For this reason, we do not present here the excellent fits achieved by the models on the fitting samples.

increases (however, perhaps, not as much as might have been thought *a priori*. See, in particular, *Economics PhD programs*). As with the simulated datasets, the SV, EW, and MR models with continuous variables outperform their binary counterparts (with one exception in 36 comparisons) but DRb dominates DRc. As to the SVc versus DEBA comparison, SVc outperforms DEBA by some margin on the *Golf rankings* and *Golf earnings* datasets but DEBA dominates SVc on the other two datasets. From a statistical viewpoint, these two datasets differ from the others in that the *Economics PhD programs* has low variability of cue validities and *Consumer reports: Digital Cameras* has low average cue inter-correlation.

Differences between models on the different datasets are highlighted in Figure 3 where we again show the performances of SVc, DEBA, EWc, DRb, and MRc. Although the datasets have some similarities (e.g., the validities of the first cue and the correlations between the criteria and the means of the predictor variables are almost the same), other differences (notably variability in cue validities and levels of cue inter-correlations) are sufficient to change the relative effectiveness of the models. SVc is very effective for the *Golf rankings* and *Golf earnings* data, EWc predicts the *Economics PhD programs* well and DEBA is best for the *Consumer reports: Digital Cameras* data. As to the two benchmark models, DRb generally has the lowest performance but, with the exception of the *Consumer reports: Digital cameras* dataset, is close to DEBA. For all datasets, MRc is always one of the better models but it does not dominate the other models. Finally, we emphasize once again the close fit between our model predictions and the empirical realizations demonstrated in Table 12. Taken as a whole, our theoretical models account for complex patterns of data.

VI. Discussion

We have mapped regions of rationality by studying a class of decisions that involve choosing the best of m ($m \geq 2$) alternatives on the basis of k ($k \geq 1$) cues or attributes. As such, this is a common task in inference and also has applications to preference (cf., Hogarth & Karelaia, in press). We have shown – through theory, simulation, and empirical demonstration – that certain simple, heuristic models can have effective performance relative to more complex, sophisticated benchmarks and, indeed, when data are scarce can, on occasion, perform better than the latter. More importantly, our theoretical analysis predicted differential model performance in a wide range of environments. Thus, for example, for our empirical datasets we predicted – and later verified – that EWc would be the best of the simple models for *Economics PhD programs* but SVc the best for *Golf rankings*.

General trends concerning relative model performance have, of course, been known for some time (e.g., effects of inter-correlations between cues or attributes, cf., Einhorn & Hogarth, 1975; Payne et al., 1993). However, the advantage of our approach is that we can specify *a priori* the *combined* effects of different environmental characteristics such as variability in cue validities, inter-correlations, level of error, and so on. Moreover, we observed that the effects of “tradeoffs” between such factors are complex and often defy simple description. The terrain that we have mapped has many dimensions.

One factor we did not consider was the effects of sampling alternatives from the underlying populations in biased or non-random ways. Clearly, results would be different if sampling excluded certain profiles of alternatives such as those likely to

dominate others or be dominated.¹⁹ On the other hand, our theoretical method allows us to make case-by-case predictions such that – through suitable aggregation – we could make predictions for samples drawn in specific, non-random ways provided the same sampling procedures are used in both fitting and holdout samples. Showing the effects of such non-random sampling is thus a straightforward task that can be addressed in future research.

This paper is also limited by the criterion used to measure model effectiveness, i.e., the emphasis on probability of correct choices. This might seem restrictive in that it assumes a “hit or miss” criterion with no consideration as to how “good” the other alternatives are. We accept this limitation in the present work but emphasize that our methodology can be easily extended to other loss functions.

First, for simplicity, consider an example of binary choice using a single variable (SV) where our methodology is used to determine the probability that alternative A is better than alternative B. That is, instead of determining the probability that Y_a is greater than Y_b (or equivalently $Y_a - Y_b > 0$), we could also consider the probability that $Y_a - Y_b > c$ where $c > 0$. In order to find this value, we need to modify the inequalities involving error differences. For example, the inequality (6), written as $\rho_{yx}(X_a - X_b) > \varepsilon_b - \varepsilon_a$, becomes:

$$\rho_{yx}(X_a - X_b) - c > \varepsilon_b - \varepsilon_a \quad (6')$$

and we proceed as before with the calculations. Moreover, by repeating this calculation for different values of c , one can investigate *how much* better Y_a is likely to be compared to Y_b . As an example, one can calculate the value of c for which $P\{Y_a - Y_b > c\} > 0.5$ or other meaningful levels of probability.

¹⁹ Note that we would not be able to apply our “overall formulas” (e.g., equation 13) to these populations because they assume unbiased, random sampling.

Second, our theoretical models can be used to specify not just the probability that one alternative will be correctly selected but also the probabilities for all alternatives. For example, imagine choosing between three alternatives A, B, and C using the SV model and having observed $x_a > x_b > x_c$. Above, we calculated the probability $P\{(Y_a > Y_b | X_a = x_a > X_b = x_b) \cap (Y_a > Y_c | X_a = x_a > X_c = x_c)\}$. However, we could also have calculated the probability that B is the largest, that is $P\{(Y_b > Y_a | X_a = x_a > X_b = x_b) \cap (Y_b > Y_c | X_b = x_b > X_c = x_c)\}$ and so on. In other words, we can specify the probabilities associated with all possibilities. Given such distributions over possible outcomes, it is straightforward to consider the effects of different loss functions, a topic we also leave for further research.

Future work could also build on our theoretical approach to consider variations of the models we have examined here. For example, models might involve mixtures of categorical and continuous variables or the effects of different types of error. How, for instance, would simple models perform when there are errors in the variables (perhaps due to measurement problems) or missing values? In addition, it will be important to investigate effects due to deviations from assumptions of normal distributions examined in this paper. Clearly many further elaborations can be undertaken.

Our work has particular implications for decision making when attention is a scarce resource. As stated by Simon (1978):

In a world in which information is relatively scarce, and where problems for decision are few and simple, information is almost always a positive good. In a world where attention is a major scarce resource, information may be an expensive luxury, for it may turn our attention from what is important to what is unimportant. We cannot afford to attend to information simply because it is there (Simon, 1978, p. 13).

By way of illustration, Simon described executives whose management information systems provide excessive, detailed information. Our work also identified regions where more information does not necessarily lead to better decisions and, if we assume that more complex models require more cognitive effort (or computational cost), there are many areas where there is no tradeoff between accuracy and effort. For example, in cases B and D illustrated in Figure 2, the simple SVc model is more accurate than the other models indicated across almost the whole range of conditions and, yet, it uses less information. On the other hand, EWc is generally best in case C where SVc lags behind the other models. However, EWc uses more information than both SVc and DEBA such that one can ask whether the additional predictive ability is worth its cost.

The models we examined might also be used in applied areas such as consumer research (cf., Bettman, Luce, & Payne, 1998). That is, instead of assuming that consumers make tradeoffs across many attributes, simpler SV or EW models can be constructed after eliciting a few simple questions concerning, say, relative importance of attributes. For example, we have shown elsewhere that if people have loose preferences characterized by binary attributes, the outputs of DEBA are remarkably consistent with more complex, linear tradeoff models (Hogarth & Karelaia, in press). However, it would be a mistake to assume that consumer preferences can always be modeled by one simple model (e.g., EW). Indeed, our theoretical analysis provides the basis for deciding which models are suited to different environments.

As suggested, our models clearly have many implications for prescriptive work. In addition to determining when heuristic models are appropriate for specific, applied problems in, for example, forecasting performance, personnel assessment and

recruitment, medical decision making, etc, heuristic models are useful supports for more complicated, decision analytical modeling. When, for instance, does one need to assess tradeoffs precisely, or can heuristic-based simplifications suffice? Here theory such as developed in this paper has great practical use for determining which heuristic to use, and when.

Finally, in a world where attention is the scarce resource, we note that “rational behavior” consists of finding the appropriate match between a decision rule and the task with which it is confronted – a principle that is valid for both descriptive and prescriptive approaches to decision making. On considering both dimensions, therefore, we do not need to assume unlimited computational capacity. However, by relaxing this assumption, we incur two costs. The first, analyzed in this paper, is to identify the task conditions under which specific heuristic rules are and are not effective, i.e., to develop maps of the regions of rationality. The second, that awaits further research, is to elucidate the conditions under which people do or do not acquire such knowledge. In other words, how do people build maps of their decision making terrain? To be effective, people do not need much computational ability to make choices in the mazes that define their environments. However, they do need task-specific knowledge or maps.

References

- Baucells, M., Carrasco, J. A., & Hogarth, R. M. (2005). *Cumulative dominance and heuristic performance in binary multiattribute choice*. Barcelona: Working paper IESE, UPC, and UPF.
- Bettman, J. R., Luce, M. F., & Payne, J. W. (1998). Constructive consumer choice processes. *Journal of Consumer Research*, 25, 187-217.
- Conlisk, J. (1996). Why bounded rationality? *Journal of Economic Literature*, 34 (2), 669-700.
- Dawes, R. M., (1979). The robust beauty of improper linear models. *American Psychologist*, 34, 571-582.
- Dawes, R. M., & Corrigan, B. (1974). Linear models in decision making. *Psychological Bulletin*, 81, 95-106.
- Dawes, R. M., Faust, D., & Meehl, P. E. (1989). Clinical versus actuarial judgment. *Science*, 243, 1668-1674.
- Einhorn, H. J., & Hogarth, R. M. (1975). Unit weighting schemes for decision making. *Organizational Behavior and Human Performance*, 13, 171-192.
- Fasolo, B., McClelland, G. H., & Todd, P. M. (in press). Escaping the tyranny of choice: When fewer attributes make choice easier. *Marketing Theory*.
- Gigerenzer, G., & Goldstein, D. G. (1996). Reasoning the fast and frugal way: Models of bounded rationality. *Psychological Review*, 103, 650-669.
- Gigerenzer, G., & Selten, R. (2001). (Eds.) *Bounded rationality: The adaptive toolbox*. Cambridge, MA: The MIT Press.
- Gigerenzer, G., Todd, P. M., and the ABC Research Group. (1999). *Simple heuristics that make us smart*. New York: Oxford University Press.

- Goldstein, D. G., & Gigerenzer, G. (2002). Models of ecological rationality: The recognition heuristic. *Psychological Review*, 109 (1), 75-90.
- Goldstein, W. M., & Hogarth, R. M. (1997). Judgment and decision research: Some historical context. In W. M. Goldstein & R. M. Hogarth (eds.), *Research on judgment and decision making: Currents, connections and controversies*. New York: Cambridge University Press.
- Hertwig, R., & Todd, P. M. (2003). More is not always better: The benefits of cognitive limits. In D. J. Hardman & L. Macchi (eds.), *Thinking: Psychological perspectives on reasoning, judgment and decision making*. Chichester, England: Wiley.
- Hogarth, R. M., & Karelaia, N. (2004). "Take-the-Best" and other simple strategies: Why and when they work "well" in binary choice. Barcelona: Universitat Pompeu Fabra, DEE working paper.
- Hogarth, R. M., & Karelaia, N. (2005). Ignoring information in binary choice with continuous variables: When is less "more"? *Journal of Mathematical Psychology*, 49, 115-124.
- Hogarth, R. M., & Karelaia, N. (in press). Simple models for multi-attribute choice with many alternatives: When it does and does not pay to face tradeoffs with binary attributes. *Management Science*.
- Kahneman, D. (2003). Maps of bounded rationality: Psychology for behavioral economics. *American Economic Review*, 93(5), 1449-1475.
- Kahneman, D., Slovic, P., & Tversky, A. (1982). (Eds.) *Judgment under uncertainty: Heuristics and biases*. New York: Cambridge University Press.
- Karelaia, N., & Hogarth, R. M. (in preparation). *On predicting performance of DEBA models in the presence of error*. Barcelona: Universitat Pompeu Fabra
- Katsikopoulos, K., & Martignon, L. (2003). *Paired comparisons: Ordering or adding*

- attributes?* Working paper, Max-Planck Institute for Human Development, Berlin.
- Kleinmuntz, B. (1990). Why we still use our heads instead of formulas: Toward an integrative approach. *Psychological Bulletin*, 107, 296-310.
- Koehler, D., & Harvey, N. (eds.) (2004). *Blackwell Handbook of Judgment and Decision Making*. Oxford, UK: Blackwell.
- Martignon, L., & Hoffrage, U. (1999). Why does one-reason decision making work? A case study in ecological rationality. In G Gigerenzer, P. M. Todd and the ABC Research Group. *Simple heuristics that make us smart* (pp. 119-140). New York: Oxford University Press.
- Martignon, L., & Hoffrage, U. (2002). Fast, frugal, and fit: Simple heuristics for paired comparison. *Theory and Decision*, 52, 29-71.
- Payne, J. W., Bettman, J. R., & Johnson, E. J. (1993). *The adaptive decision maker*. New York: Cambridge University Press.
- Shanteau, J., & Thomas, R. P. (2000). Fast and frugal heuristics: What about unfriendly environments? *Behavioral and Brain Sciences*, 23, 762-763.
- Simon, H. A. (1956). Rational choice and the structure of environments. *Psychological Review*, 63, 129-138.
- Simon, H. A. (1978). Rationality as process and product of thought. *American Economic Review*, 68 (2), 1-16
- Simon, H. A. (1991). *Models of my life*. New York: Basic Books.
- Thorngate, W. (1980). Efficient decision heuristics. *Behavioral Science*, 25, 219-225.
- Tversky, A. (1972). Elimination by aspects: A theory of choice. *Psychological Review*, 79, 281-299.

Table 1 -- Models tested

<u>(A). Single Variable (SV) models</u>		<u>Prior information*</u>	<u>Information to consult</u>	<u>Calculations</u>	<u>Number of comparisons</u>	
					<u>min</u>	<u>max</u>
1	Lexicographic -- SVC	Choice depends solely on cue with the greatest validity (see, e.g., Payne et al., 1993).				
2	Lexicographic -- SVb	Model 1 above but based on binary variables.	Rank-order of importance of cues	None	$(m - 1)$	$\frac{m(m - 1)}{2}$
3	DEBA	Deterministic version of Tversky's (1972) elimination-by-aspects model (EBA). For binary choice, this is the same as the take-the-best model of Gigerenzer and Goldstein (1996).				Variable
<u>(B). Equal weight (EW) models</u>						
4	EWc	All cues are accorded equal weight (see, e.g., Einhorn & Hogarth, 1975).	None	All	m sums	$(m - 1)$
5	EWb	All cues are accorded equal weight (see, e.g., Dawes, 1979),				
<u>(C). Hybrid models</u>						
6	EW/DEBA	Choose according to equal weights. If this results in a tie, use DEBA to resolve the choice (Hogarth & Karelaia, in press).	Rank-order of importance of cues	All	m sums	$(m - 1)$
7	EW/SVb	Choose according to equal weights. If this results in a tie, resolve conflict by the single most important variable				
<u>(D). Domran (DR) models</u>						
8	DRc	If an alternative dominates, choose it. Otherwise, choose at random between non-dominated alternatives.	None	All	None	$k(m - 1)$
9	DRb	Same as DRc except based on binary variables.				
<u>(E). Multiple regression (MR)</u>						
10	MRc	Well-known statistical model.	Importance of cues	All	km products, m sums	$(m - 1)$
11	MRb	Same as MRc except based on binary variables.				

* For all models, the decision maker is assumed to know the sign of the zero order correlation between cues and the criterion.

NOTE: m = number of alternatives, k = number of attributes, cues

Table 2 – Key formulas for different models using continuous variables (for choosing best of three)

	<u>Accuracy conditions</u>	<u>Error variances</u>	<u>Upper integration limits,</u> l_{ab} and l_{ac}
<u>Single variable (SVc)</u>			
$Y_a = \rho_{yx} X_a + \varepsilon_a$	$\rho_{yx} (X_a - X_b) > \varepsilon_b - \varepsilon_a$	$2(1 - \rho_{yx}^2)$	$\frac{\rho_{yx} (x_a - x_b)}{\sqrt{2(1 - \rho_{yx}^2)}}$
$Y_b = \rho_{yx} X_b + \varepsilon_b$	$\rho_{yx} (X_a - X_c) > \varepsilon_c - \varepsilon_a$		$\frac{\rho_{yx} (x_a - x_c)}{\sqrt{2(1 - \rho_{yx}^2)}}$
$Y_c = \rho_{yx} X_c + \varepsilon_c$			
<u>Equal weights (EWC)</u>			
$Y_a = (\rho_{y\bar{x}} / \sigma_{\bar{x}}) \bar{X}_a + v_a$	$(\rho_{y\bar{x}} / \sigma_{\bar{x}}) (\bar{X}_a - \bar{X}_b) > v_b - v_a$	$2(1 - \rho_{y\bar{x}}^2)$	$\frac{\rho_{y\bar{x}} (\bar{x}_a - \bar{x}_b)}{\sigma_{\bar{x}} \sqrt{2(1 - \rho_{y\bar{x}}^2)}}$
$Y_b = (\rho_{y\bar{x}} / \sigma_{\bar{x}}) \bar{X}_b + v_b$	$(\rho_{y\bar{x}} / \sigma_{\bar{x}}) (\bar{X}_a - \bar{X}_c) > v_c - v_a$		$\frac{\rho_{y\bar{x}} (\bar{x}_a - \bar{x}_c)}{\sigma_{\bar{x}} \sqrt{2(1 - \rho_{y\bar{x}}^2)}}$
$Y_c = (\rho_{y\bar{x}} / \sigma_{\bar{x}}) \bar{X}_c + v_c$			
<u>Multiple regression (MRc)</u>			
$Y_a = \hat{Y}_a + u_a$	$(\hat{Y}_a - \hat{Y}_b) > u_b - u_a$	$2(1 - R_{adj}^2)$	$\frac{(\hat{y}_a - \hat{y}_b)}{\sqrt{2(1 - R_{adj}^2)}}$
$Y_b = \hat{Y}_b + u_b$	$(\hat{Y}_a - \hat{Y}_c) > u_c - u_a$		
$Y_c = \hat{Y}_c + u_c$			$\frac{(\hat{y}_a - \hat{y}_c)}{\sqrt{2(1 - R_{adj}^2)}}$

Notes:

$$1. \quad \rho_{y\bar{x}} = \bar{\rho}_{yx} \sqrt{\frac{k}{1 + (k-1)\bar{\rho}_{x_i x_j}}} \quad \text{where } k = \text{number of } x \text{ variables, } \bar{\rho}_{yx} = \text{average correlation between } y \text{ and the } x\text{'s, and}$$

$\bar{\rho}_{x_i x_j} = \text{average inter-correlations amongst the } x\text{'s.}$

$$2. \quad \sigma_{\bar{x}} = \sqrt{\frac{1}{k} \left(1 + (k-1)\bar{\rho}_{x_i x_j} \right)}$$

$$3. \quad R_{adj}^2 = 1 - (1 - R^2) \frac{(n-1)}{(n-k)} \quad \text{where } n = \text{number of observations.}$$

4. The probability of optimal choice out of three is given by:

$$\int_{-\infty}^{l_{ab}} \int_{-\infty}^{l_{ac}} \frac{1}{\pi\sqrt{3}} e^{-\frac{2}{3}(z_1^2 - z_1 z_2 + z_2^2)} dz_1 dz_2,$$

with upper integration limits being specific for different choice strategies.

$$5. \quad \text{Correlations between the standardized normal variables } z_1 = \frac{\mathcal{E}_b - \mathcal{E}_a}{\sqrt{2(1 - \rho_{yx}^2)}} \quad \text{and } z_2 = \frac{\mathcal{E}_c - \mathcal{E}_a}{\sqrt{2(1 - \rho_{yx}^2)}} \quad \text{are all equal to } \frac{1}{2} \text{ - see Appendix A.}$$

Table 3 – Formulas for overall predictive accuracy of models using continuous variables
(for choosing best of m)

<u>Model</u>	<u>V_d</u>	<u>Upper limits of integration</u> d_i^* , for $i = \overline{1, m-1}$
<u>SVc</u>	$\begin{pmatrix} 2 & \dots & 1 \\ \dots & \dots & \dots \\ 1 & \dots & 2 \end{pmatrix}$	$\frac{\rho_{yx} d_i}{\sqrt{2(1 - \rho_{yx}^2)}}$
<u>EWc</u>	$\begin{pmatrix} \sqrt{2}\sigma_{\bar{x}} & \dots & \sigma_{\bar{x}}^2 \\ \dots & \dots & \dots \\ \sigma_{\bar{x}}^2 & \dots & \sqrt{2}\sigma_{\bar{x}} \end{pmatrix}$	$\frac{\rho_{y\bar{x}} d_i}{\sigma_{\bar{x}} \sqrt{2(1 - \rho_{y\bar{x}}^2)}}$
<u>MRC</u>	$\begin{pmatrix} \sqrt{2R_{adj}^2} & \dots & R_{adj}^2 \\ \dots & \dots & \dots \\ R_{adj}^2 & \dots & \sqrt{2R_{adj}^2} \end{pmatrix}$	$\frac{d_i}{\sqrt{2(1 - R_{adj}^2)}}$

Notes:

Overall expected probability of success is given by:

$$m \int_0^\infty \dots \int_0^\infty \varphi(d | \mu_d, V_d) \left[\int_{-\infty}^{d_1^*} \dots \int_{-\infty}^{d_{m-1}^*} \varphi(z | \mu_z, V_z) dz_1 \dots dz_{m-1} \right] dd_1 \dots dd_{m-1}$$

where $z' = (z_1, z_2, \dots, z_{m-1})$, $d' = (d_1, d_2, \dots, d_{m-1})$, and $V_z = \begin{pmatrix} 1 & \dots & 1/2 \\ \dots & \dots & \dots \\ 1/2 & \dots & 1 \end{pmatrix}$.

V_d and d_i^* , for $i = \overline{1, m-1}$, are specific for different choice strategies.

See also notes to Table 2.

Table 4 – Key formulas for different models using binary variables (for choosing best of three)

<u>Model</u>		<u>Upper integration limits</u>	
		h_{ab}	h_{ac}
<u>Single variable (SVb)</u>	$Y = a_{SVb} + \frac{\rho_{yw}}{\sigma_w} W + \zeta$	$\frac{\rho_{yw}(w_a - w_b)}{\sigma_w \sqrt{2(1 - \rho_{yw}^2)}}$	$\frac{\rho_{yw}(w_a - w_c)}{\sigma_w \sqrt{2(1 - \rho_{yw}^2)}}$
<u>Equal weights (EWb)</u>	$Y = a_{EWb} + \frac{\rho_{y\bar{w}}}{\sigma_{\bar{w}}} \bar{W} + \xi$	$\frac{\rho_{y\bar{w}}(\bar{w}_a - \bar{w}_b)}{\sigma_{\bar{w}} \sqrt{2(1 - \rho_{y\bar{w}}^2)}}$	$\frac{\rho_{y\bar{w}}(\bar{w}_a - \bar{w}_c)}{\sigma_{\bar{w}} \sqrt{2(1 - \rho_{y\bar{w}}^2)}}$
<u>Multiple regression (MRb)</u>	$Y = \hat{Y} + \zeta$	$\frac{(\hat{y}_a - \hat{y}_b)}{\sqrt{2(1 - R_{adj}^2)}}$	$\frac{(\hat{y}_a - \hat{y}_c)}{\sqrt{2(1 - R_{adj}^2)}}$

(\hat{Y} and R_{adj}^2 are based on binary variables W 's)

Notes:

$$1. \quad \rho_{y\bar{w}} = \bar{\rho}_{yw} \sqrt{\frac{k}{1 + (k-1)\bar{\rho}_{w_i w_j}}},$$

where k = number of x variables,

$\bar{\rho}_{yw}$ = average correlation between y and the w 's, and

$\bar{\rho}_{w_i w_j}$ = average inter-correlations amongst the w 's.

$$2. \quad \sigma_{\bar{w}} = \sqrt{\frac{1}{k} (1 + (k-1)\bar{\rho}_{w_i w_j})}$$

$$3. \quad R_{adj}^2 = 1 - \frac{(1 - R^2)(n-1)}{(n-k)}$$

where n = number of observations.

4. The probability of optimal choice out of three is given by

$$\int_{-\infty}^{h_{ab}} \int_{-\infty}^{h_{ac}} \frac{1}{\pi\sqrt{3}} e^{-\frac{2}{3}(z_1^2 - z_1 z_2 + z_2^2)} dz_1 dz_2,$$

with upper integration limits being specific for different choice strategies.

5. Since the error terms have means of zero, the intercepts a_{SVb} and a_{EWb} are equal

to $-\rho_{yw}$ and $-\frac{\rho_{y\bar{w}}}{2\sigma_{\bar{w}}}$, respectively.

Table 5 -- Experimental design

subcases	<u>Case A (k = 3)</u>					<u>Case B (k = 3)</u>					<u>Case C (k = 5)</u>					<u>Case D (k = 5)</u>				
	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>
Variability of cue validities: max - min	0.30	0.39	0.49	0.60	0.70	0.29	0.39	0.50	0.59	0.69	0.14	0.25	0.30	0.34	0.44	0.00	0.10	0.30	0.40	0.50
Validity of the first cue	0.39	0.49	0.59	0.69	0.79	0.39	0.49	0.59	0.69	0.79	0.34	0.44	0.49	0.54	0.64	0.29	0.39	0.59	0.69	0.79
Average validity	0.26	0.29	0.33	0.36	0.39	0.26	0.29	0.32	0.36	0.39	0.25	0.29	0.34	0.38	0.44	0.29	0.33	0.37	0.40	0.43
Average intercorrelation of x's	0.00	0.00	0.00	0.00	0.00	0.49	0.49	0.49	0.49	0.49	0.10	0.10	0.10	0.10	0.10	0.60	0.60	0.60	0.60	0.60
Correlation: y and \bar{x}	0.45	0.51	0.57	0.63	0.68	0.31	0.36	0.39	0.44	0.48	0.47	0.56	0.65	0.73	0.84	0.35	0.40	0.45	0.48	0.52
R ² (MR) -- fit	0.35	0.43	0.52	0.63	0.76	0.31	0.39	0.49	0.62	0.78	0.42	0.51	0.61	0.70	0.88	0.34	0.38	0.51	0.63	0.78
(n-1)/(n-k)	1.12	1.12	1.12	1.12	1.12	1.12	1.12	1.12	1.12	1.12	1.27	1.27	1.27	1.27	1.27	1.27	1.27	1.27	1.27	1.27

Table 6 -- Predictions and realizations (% correct) for simulated data: Best of "two"

		<u>Case A</u>					<u>Case B</u>					<u>Case C</u>					<u>Case D</u>					<u>Mean realizations</u>	
<u>subcases</u>		<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>		
<u>Single Variable models</u>																							
1	Lexicographic -- SVc	Predictions	63	67	71	75	80	63	67	71	74	79	61	65	67	69	72	60	63	70	75	80	69
		<u>Realizations</u>	<u>62</u>	<u>66</u>	70	74	80	63	66	71	75	80	<u>59</u>	<u>65</u>	<u>66</u>	<u>68</u>	<u>72</u>	<u>59</u>	<u>62</u>	71	75	79	
2	Lexicographic -- SVb	Predictions	59	61	64	67	69	59	61	64	67	69	58	60	61	63	65	57	59	64	66	69	63
		<u>Realizations</u>	<u>60</u>	<u>62</u>	<u>64</u>	<u>66</u>	<u>69</u>	<u>59</u>	<u>61</u>	<u>64</u>	<u>67</u>	<u>69</u>	<u>58</u>	<u>59</u>	<u>62</u>	<u>63</u>	<u>65</u>	<u>57</u>	<u>60</u>	<u>64</u>	<u>67</u>	<u>69</u>	
3	DEBA	Predictions	63	66	68	72	74	61	62	64	66	68	63	66	69	72	77	60	63	66	67	70	67
		<u>Realizations</u>	<u>64</u>	67	<u>69</u>	<u>72</u>	<u>74</u>	<u>62</u>	<u>62</u>	<u>64</u>	<u>66</u>	<u>68</u>	<u>63</u>	<u>67</u>	<u>69</u>	<u>73</u>	<u>77</u>	<u>60</u>	63	<u>66</u>	<u>68</u>	<u>70</u>	
<u>Equal Weight models</u>																							
4	EWc	Predictions	66	68	70	72	75	61	62	63	65	66	66	69	73	77	83	62	63	66	67	68	68
		<u>Realizations</u>	65	67	<u>69</u>	<u>71</u>	<u>73</u>	<u>61</u>	<u>61</u>	<u>63</u>	<u>65</u>	<u>66</u>	65	69	73	76	83	<u>62</u>	63	<u>66</u>	<u>67</u>	<u>69</u>	
5	EWb	Predictions	62	63	65	67	68	59	60	62	63	64	63	66	69	71	75	61	62	64	65	66	65
		<u>Realizations</u>	<u>61</u>	<u>63</u>	<u>64</u>	<u>67</u>	<u>68</u>	<u>58</u>	<u>60</u>	<u>61</u>	<u>63</u>	<u>63</u>	<u>63</u>	<u>65</u>	<u>68</u>	<u>71</u>	<u>75</u>	63	63	<u>64</u>	<u>66</u>	<u>66</u>	
<u>Hybrid models</u>																							
6	EW/DEBA	Predictions	63	65	68	70	73	60	62	63	65	67	64	67	71	73	78	61	63	64	66	68	67
		<u>Realizations</u>	<u>64</u>	<u>65</u>	<u>69</u>	<u>71</u>	<u>73</u>	<u>60</u>	<u>62</u>	<u>64</u>	<u>66</u>	<u>67</u>	<u>63</u>	<u>67</u>	<u>70</u>	<u>73</u>	<u>78</u>	<u>60</u>	63	<u>65</u>	<u>67</u>	<u>68</u>	
7	EW/SVb	Predictions	63	65	67	70	72	60	61	63	65	66	64	67	70	73	77	61	63	64	66	68	66
		<u>Realizations</u>	<u>63</u>	<u>64</u>	<u>68</u>	<u>70</u>	<u>72</u>	<u>59</u>	<u>62</u>	<u>63</u>	<u>65</u>	<u>66</u>	<u>63</u>	<u>66</u>	<u>70</u>	<u>72</u>	<u>77</u>	<u>60</u>	63	<u>65</u>	<u>66</u>	<u>67</u>	
<u>Domran models</u>																							
8	DRc	Fit	57	57	58	59	60	58	59	59	62	63	54	54	54	55	56	58	59	61	61	62	58
		<u>Realizations</u>	<u>57</u>	<u>57</u>	<u>58</u>	<u>59</u>	<u>60</u>	<u>58</u>	<u>59</u>	<u>60</u>	<u>60</u>	<u>64</u>	<u>54</u>	<u>54</u>	<u>54</u>	<u>55</u>	<u>55</u>	<u>57</u>	<u>59</u>	<u>61</u>	<u>61</u>	<u>62</u>	
9	DRb	Fit	61	62	63	64	67	59	60	60	63	64	60	61	62	65	67	60	62	63	64	65	63
		<u>Realizations</u>	<u>60</u>	<u>62</u>	<u>63</u>	<u>64</u>	<u>66</u>	<u>59</u>	<u>60</u>	<u>62</u>	<u>63</u>	<u>64</u>	<u>59</u>	<u>61</u>	<u>63</u>	<u>65</u>	<u>68</u>	<u>60</u>	<u>61</u>	<u>63</u>	<u>64</u>	<u>65</u>	
<u>Multiple regression</u>																							
10	MRc	Predictions	69	72	75	79	83	68	70	74	78	84	72	74	77	80	88	69	70	74	78	84	70
		<u>Realizations</u>	<u>64</u>	67	71	76	82	<u>62</u>	<u>65</u>	71	76	82	<u>61</u>	<u>67</u>	<u>70</u>	76	85	<u>56</u>	<u>59</u>	<u>67</u>	<u>72</u>	80	
11	MRb	Predictions	62	62	64	65	67	63	63	65	67	69	65	66	66	67	67	65	65	67	68	70	60
		<u>Realizations</u>	<u>56</u>	<u>58</u>	<u>61</u>	<u>64</u>	<u>66</u>	<u>56</u>	<u>59</u>	<u>62</u>	<u>66</u>	<u>70</u>	<u>53</u>	<u>56</u>	<u>58</u>	<u>59</u>	<u>61</u>	<u>54</u>	<u>54</u>	<u>57</u>	<u>61</u>	<u>66</u>	
<u>Mean realizations</u>		<u>61</u>	<u>63</u>	<u>66</u>	<u>69</u>	<u>71</u>	<u>60</u>	<u>62</u>	<u>64</u>	<u>67</u>	<u>69</u>	<u>60</u>	<u>63</u>	<u>66</u>	<u>68</u>	<u>72</u>	<u>59</u>	<u>61</u>	<u>64</u>	<u>67</u>	<u>69</u>		

Note: **Bold** figures denote largest realization in each column or second largest if MRc is largest.

Table 7 -- Predictions and realizations (% correct) for simulated data: Best of "three"

		<u>Case A</u>					<u>Case B</u>					<u>Case C</u>					<u>Case D</u>					<u>Mean</u>	
<u>subcases</u>		<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>realizations</u>	
<u>Single Variable models</u>																							
1	Lexicographic -- SVC	Predictions	49	54	59	64	71	49	54	59	64	71	47	51	54	56	61	45	49	58	64	70	
	<u>Realizations</u>	<u>50</u>	<u>54</u>	58	64	71	51	53	58	64	70	<u>47</u>	<u>52</u>	<u>53</u>	<u>55</u>	<u>62</u>	<u>46</u>	<u>49</u>	58	64	70	<u>57</u>	
2	Lexicographic -- SVb	Predictions	44	48	51	53	56	44	47	50	53	57	43	46	47	49	52	41	44	50	53	57	
	<u>Realizations</u>	<u>44</u>	<u>47</u>	<u>50</u>	<u>53</u>	<u>57</u>	<u>44</u>	<u>47</u>	<u>50</u>	<u>52</u>	<u>56</u>	<u>43</u>	<u>44</u>	<u>47</u>	<u>48</u>	<u>52</u>	<u>41</u>	<u>44</u>	<u>51</u>	<u>54</u>	<u>57</u>	<u>49</u>	
3	DEBA	Predictions	50	53	56	60	64	46	49	51	54	56	49	53	57	60	67	45	49	52	55	58	
	<u>Realizations</u>	<u>50</u>	<u>54</u>	<u>56</u>	<u>59</u>	<u>63</u>	<u>46</u>	<u>48</u>	<u>50</u>	<u>53</u>	<u>55</u>	<u>48</u>	<u>53</u>	<u>57</u>	<u>61</u>	<u>66</u>	<u>45</u>	<u>49</u>	<u>53</u>	<u>55</u>	<u>57</u>	<u>54</u>	
<u>Equal Weight models</u>																							
4	EWc	Predictions	52	55	58	61	64	46	48	50	51	53	53	57	62	67	75	48	50	52	53	55	
	<u>Realizations</u>	52	55	<u>57</u>	<u>61</u>	<u>64</u>	<u>45</u>	<u>48</u>	<u>51</u>	<u>51</u>	<u>52</u>	52	57	62	66	74	<u>47</u>	50	<u>51</u>	<u>52</u>	<u>53</u>	<u>55</u>	
5	EWb	Predictions	47	49	51	53	55	44	46	47	49	50	49	52	56	59	65	46	48	50	51	53	
	<u>Realizations</u>	<u>47</u>	<u>49</u>	<u>51</u>	<u>54</u>	<u>55</u>	<u>44</u>	<u>45</u>	<u>47</u>	<u>47</u>	<u>50</u>	<u>48</u>	<u>51</u>	<u>56</u>	<u>58</u>	<u>65</u>	<u>46</u>	<u>47</u>	<u>52</u>	<u>51</u>	<u>54</u>	<u>51</u>	
<u>Hybrid models</u>																							
6	EW/DEBA	Predictions	49	52	55	58	61	46	47	49	52	54	50	54	58	62	68	46	48	51	52	55	
	<u>Realizations</u>	<u>50</u>	<u>53</u>	<u>54</u>	<u>58</u>	<u>62</u>	<u>46</u>	<u>47</u>	<u>49</u>	<u>52</u>	<u>53</u>	<u>50</u>	<u>54</u>	<u>58</u>	<u>61</u>	<u>68</u>	<u>47</u>	<u>49</u>	<u>50</u>	<u>51</u>	<u>54</u>	<u>53</u>	
7	EW/SVb	Predictions	49	51	54	57	60	45	47	49	51	53	50	54	57	61	67	46	48	51	52	55	
	<u>Realizations</u>	<u>49</u>	<u>52</u>	<u>53</u>	<u>57</u>	<u>61</u>	<u>45</u>	<u>47</u>	<u>48</u>	<u>51</u>	<u>52</u>	<u>50</u>	<u>54</u>	<u>57</u>	<u>60</u>	<u>66</u>	<u>47</u>	<u>49</u>	<u>50</u>	<u>51</u>	<u>54</u>	<u>53</u>	
<u>Domran models</u>																							
8	DRc	Fit	44	44	45	45	47	48	48	50	51	52	38	39	39	39	39	47	48	49	49	50	
	<u>Realizations</u>	<u>44</u>	<u>44</u>	<u>46</u>	<u>45</u>	<u>47</u>	<u>48</u>	<u>48</u>	<u>49</u>	<u>51</u>	<u>52</u>	<u>38</u>	<u>39</u>	<u>39</u>	<u>39</u>	<u>39</u>	<u>46</u>	<u>48</u>	<u>48</u>	<u>49</u>	<u>51</u>	<u>46</u>	
9	DRb	Fit	50	51	52	54	56	47	49	50	51	53	49	50	52	53	57	50	51	52	52	54	
	<u>Realizations</u>	<u>49</u>	<u>51</u>	<u>54</u>	<u>54</u>	<u>55</u>	<u>48</u>	<u>49</u>	<u>51</u>	<u>49</u>	<u>53</u>	<u>49</u>	<u>50</u>	<u>52</u>	<u>54</u>	<u>55</u>	48	50	<u>51</u>	<u>52</u>	<u>54</u>	<u>51</u>	
<u>Multiple regression</u>																							
10	MRc	Predictions	57	60	64	69	76	52	55	59	63	70	58	63	67	71	82	52	53	58	62	68	
	<u>Realizations</u>	<u>51</u>	57	60	67	74	<u>49</u>	<u>52</u>	<u>57</u>	<u>63</u>	<u>69</u>	<u>49</u>	<u>55</u>	62	67	<u>69</u>	<u>46</u>	<u>49</u>	<u>56</u>	<u>63</u>	<u>69</u>	<u>59</u>	
11	MRb	Predictions	46	47	49	51	53	47	49	51	53	56	52	52	53	53	54	52	53	55	56	58	
	<u>Realizations</u>	<u>42</u>	<u>45</u>	<u>48</u>	<u>52</u>	<u>55</u>	<u>42</u>	<u>46</u>	<u>48</u>	<u>52</u>	<u>58</u>	<u>40</u>	<u>43</u>	<u>44</u>	<u>47</u>	<u>51</u>	<u>39</u>	<u>40</u>	<u>46</u>	<u>50</u>	<u>55</u>	<u>47</u>	
<u>Mean realizations</u>			<u>48</u>	<u>51</u>	<u>53</u>	<u>57</u>	<u>60</u>	<u>46</u>	<u>48</u>	<u>51</u>	<u>53</u>	<u>56</u>	<u>47</u>	<u>50</u>	<u>53</u>	<u>56</u>	<u>47</u>	<u>48</u>	<u>52</u>	<u>54</u>	<u>57</u>		

Note: **Bold** figures denote largest realization in each column or second largest if MRc is largest.

Table 8 -- Predictions and realizations (% correct) for simulated data: Best of "four"

		<u>Case A</u>					<u>Case B</u>					<u>Case C</u>					<u>Case D</u>					<u>Mean realizations</u>
<u>subcases</u>		<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>	
<u>Single Variable models</u>																						
1 Lexicographic -- SVC	Predictions	42	46	52	58	65	41	46	52	58	65	39	44	46	49	54	37	41	52	58	65	<u>50</u>
	<u>Realizations</u>	<u>41</u>	<u>46</u>	52	56	65	42	45	52	58	66	<u>38</u>	<u>44</u>	<u>45</u>	<u>49</u>	<u>55</u>	<u>37</u>	<u>41</u>	52	58	65	
2 Lexicographic -- SVb	Predictions	35	38	41	44	47	35	38	41	44	47	34	37	38	40	43	33	35	41	44	48	<u>40</u>
	<u>Realizations</u>	<u>35</u>	<u>38</u>	<u>40</u>	<u>44</u>	<u>48</u>	<u>36</u>	<u>38</u>	<u>41</u>	<u>44</u>	<u>48</u>	<u>34</u>	<u>37</u>	<u>38</u>	<u>41</u>	<u>43</u>	<u>33</u>	<u>34</u>	<u>41</u>	<u>44</u>	<u>48</u>	
3 DEBA	Predictions	41	44	47	52	56	38	39	42	45	47	41	45	50	53	60	37	39	44	45	49	<u>46</u>
	<u>Realizations</u>	<u>42</u>	<u>45</u>	<u>47</u>	<u>51</u>	<u>55</u>	<u>39</u>	<u>39</u>	<u>42</u>	<u>44</u>	<u>46</u>	<u>41</u>	<u>46</u>	<u>50</u>	<u>54</u>	<u>60</u>	<u>38</u>	<u>40</u>	<u>44</u>	<u>45</u>	<u>49</u>	
<u>Equal Weight models</u>																						
4 EWc	Predictions	45	48	51	54	58	38	40	42	44	46	45	50	56	61	70	40	42	44	46	48	<u>48</u>
	<u>Realizations</u>	45	47	<u>50</u>	<u>53</u>	<u>56</u>	<u>38</u>	<u>39</u>	<u>42</u>	<u>43</u>	<u>45</u>	44	50	54	60	70	<u>39</u>	<u>42</u>	<u>44</u>	<u>44</u>	<u>47</u>	
5 EWb	Predictions	39	41	43	46	48	36	37	39	41	42	41	45	49	53	58	38	40	42	43	45	<u>43</u>
	<u>Realizations</u>	<u>38</u>	<u>41</u>	<u>41</u>	<u>45</u>	<u>48</u>	<u>37</u>	<u>37</u>	<u>38</u>	<u>40</u>	<u>42</u>	<u>42</u>	<u>45</u>	<u>48</u>	<u>52</u>	<u>58</u>	<u>37</u>	<u>39</u>	<u>42</u>	<u>42</u>	<u>45</u>	
<u>Hybrid models</u>																						
6 EW/DEBA	Predictions	41	44	47	50	54	36	39	40	42	44	43	47	51	56	62	38	40	43	44	47	<u>45</u>
	<u>Realizations</u>	<u>41</u>	<u>44</u>	<u>46</u>	<u>51</u>	<u>54</u>	<u>38</u>	<u>39</u>	<u>42</u>	<u>42</u>	<u>44</u>	<u>43</u>	<u>47</u>	<u>51</u>	<u>55</u>	<u>61</u>	<u>37</u>	<u>41</u>	<u>42</u>	<u>42</u>	<u>47</u>	
7 EW/SVb	Predictions	40	43	46	49	53	36	38	40	42	44	42	46	50	54	60	38	40	42	43	46	<u>44</u>
	<u>Realizations</u>	<u>40</u>	<u>43</u>	<u>45</u>	<u>49</u>	<u>52</u>	<u>37</u>	<u>38</u>	<u>41</u>	<u>41</u>	<u>43</u>	<u>42</u>	<u>46</u>	<u>50</u>	<u>54</u>	<u>59</u>	<u>37</u>	<u>41</u>	<u>42</u>	<u>42</u>	<u>46</u>	
<u>Domran models</u>																						
8 DRc	Fit	37	37	38	38	39	43	44	44	45	47	31	31	31	31	31	42	42	43	43	44	<u>39</u>
	<u>Realizations</u>	<u>37</u>	<u>37</u>	<u>38</u>	<u>38</u>	<u>39</u>	42	<u>43</u>	<u>44</u>	<u>45</u>	<u>47</u>	<u>30</u>	<u>30</u>	<u>31</u>	<u>31</u>	<u>31</u>	<u>41</u>	<u>42</u>	<u>43</u>	<u>43</u>	<u>44</u>	
9 DRb	Fit	45	45	47	48	49	42	44	44	45	48	44	45	47	49	51	44	45	47	46	47	<u>46</u>
	<u>Realizations</u>	<u>43</u>	<u>45</u>	<u>47</u>	<u>48</u>	<u>50</u>	42	<u>43</u>	<u>44</u>	<u>45</u>	<u>46</u>	44	<u>45</u>	<u>47</u>	<u>48</u>	<u>50</u>	44	45	<u>46</u>	<u>47</u>	<u>48</u>	
<u>Multiple regression</u>																						
10 MRc	Predictions	50	54	58	63	71	44	47	52	57	64	51	55	61	66	78	44	46	51	56	63	<u>53</u>
	<u>Realizations</u>	45	48	54	60	69	<u>40</u>	<u>44</u>	<u>51</u>	<u>57</u>	<u>65</u>	<u>41</u>	<u>48</u>	55	61	76	<u>38</u>	<u>40</u>	<u>50</u>	<u>56</u>	65	
11 MRb	Predictions	38	39	41	43	45	39	41	43	45	49	45	45	46	46	48	45	45	48	50	52	<u>38</u>
	<u>Realizations</u>	<u>33</u>	<u>36</u>	<u>39</u>	<u>42</u>	<u>45</u>	<u>34</u>	<u>37</u>	<u>40</u>	<u>45</u>	<u>50</u>	<u>31</u>	<u>35</u>	<u>35</u>	<u>37</u>	<u>43</u>	<u>30</u>	<u>30</u>	<u>36</u>	<u>39</u>	<u>47</u>	
<u>Mean realizations</u>		<u>40</u>	<u>43</u>	<u>45</u>	<u>49</u>	<u>53</u>	<u>39</u>	<u>40</u>	<u>43</u>	<u>46</u>	<u>49</u>	<u>39</u>	<u>43</u>	<u>46</u>	<u>49</u>	<u>55</u>	<u>37</u>	<u>40</u>	<u>44</u>	<u>46</u>	<u>50</u>	

Note: **Bold** figures denote largest realization in each column or second largest if MRc is largest.

Table 9 -- Mapping relative success of different models: Realizations relative to MRc*

subcases:**		1	2	3	4	5			1	2	3	4	5		
No/low redundancy						Moderate/high redundancy									
<u>Case A</u>	SVc	1	2	1	3	4	<u>Case B</u>	SVc	-2	-1	-1	-1	0		
	DEBA	1	2	4	8	11		DEBA	3	5	7	10	15	3 cues/	
	EWc	-1	1	3	6	10		EWc	4	5	7	11	18	attributes	
	DRb	2	6	6	13	19		DRb	1	3	6	14	16		
<u>Case C</u>	SVc	1	3	9	11	17	<u>Case D</u>	SVc	0	0	-3	-2	-1		
	DEBA	1	2	5	6	14		DEBA	1	0	3	8	12	5 cues/	
	EWc	-3	-2	0	1	5		EWc	-1	-1	5	10	16	attributes	
	DRb	0	5	10	13	24		DRb	-2	-2	5	11	15		
						High noise							Low noise		

* Entries are performance (best of "three") of MRc less specified models.

Shaded regions indicate areas where MRc's advantage > 4%; bold type shows no advantage or disadvantage of MRc.

** Parameters specifying the sub-cases are provided in Table 5 -- descriptions in **bold**.

Table 10 -- Regressions of model performance (realizations) on environmental characteristics

<u>Models:</u>	<u>SVc</u>	<u>SVb</u>	<u>DEBA</u>	<u>EWc</u>	<u>EWb</u>	<u>EW/ DEBA</u>	<u>EW/ SVb</u>	<u>DRc</u>	<u>DRb</u>	<u>MRc</u>	<u>MRb</u>
<u>Regression coefficients* for:</u>											
Constant	42	47	44	42	42	43	43	65	51	43	49
Dummy1	-12	-14	-13	-13	-14	-14	-13	-13	-11	-11	-13
Dummy2	-7	-9	-8	-7	-8	-8	-8	-7	-6	-7	-9
Number of cues					1			-5		-3	
Validity of the first cue	50	29	11					50			
Correlation y and \bar{x}			32	50	37	45	32			26	-15
Variability of cue validities: max - min								-42			17
Average cue validity							19		38		
Average intercorrelation of x's					4	3		9	-4	6	
R ²	42	22
<u>Regression statistics:</u>											
R ²	0.99	0.99	0.99	0.98	0.99	0.99	0.99	0.98	0.99	0.98	0.99
Estimated standard error	1.30	0.72	1.13	1.43	1.04	1.18	1.23	1.47	0.62	1.54	0.93

* All regression coefficients are statistically significant (p < .001).

Table 11 -- Characteristics of empirical datasets

	<u>Golf rankings</u>	<u>Golf earnings</u>	<u>Economics PhD programs</u>	<u>Consumer reports: Digital cameras</u>
Variability of cue validities: max - min	0.23	0.29	0.07	0.38
Validity of the first cue	0.78	0.86	0.81	0.79
Average intercorrelation of x's	0.46	0.46	0.60	0.20
Correlation y and \bar{x}	0.78	0.84	0.89	0.80
R ² (MR) -- fit	0.68	0.81	0.81	0.73
(n-1)/(n-k)	1.07	1.07	1.04	1.09

Table 12 -- Predictions and realizations (% correct) for empirical datasets

		<u>Golf rankings</u>			<u>Golf earnings</u>			<u>PhD programs</u>			<u>Digital cameras</u>			<u>Mean realizations</u>	
		<u>Best of</u>			<u>Best of</u>			<u>Best of</u>			<u>Best of</u>				
		<u>Two</u>	<u>Three</u>	<u>Four</u>	<u>Two</u>	<u>Three</u>	<u>Four</u>	<u>Two</u>	<u>Three</u>	<u>Four</u>	<u>Two</u>	<u>Three</u>	<u>Four</u>		
<u>Single Variable models</u>															
1	Lexicographic -- SVC	Predictions	79	69	62	82	77	74	77	71	69	74	64	59	
		<u>Realizations</u>	79	72	67	81	76	73	<u>78</u>	<u>75</u>	<u>74</u>	<u>72</u>	<u>66</u>	<u>59</u>	<u>73</u>
2	Lexicographic -- SVb	Predictions	69	55	45	69	57	48	71	63	58	69	60	55	
		<u>Realizations</u>	<u>69</u>	<u>56</u>	<u>46</u>	<u>73</u>	<u>61</u>	<u>53</u>	<u>71</u>	<u>63</u>	<u>57</u>	<u>69</u>	<u>64</u>	<u>58</u>	<u>62</u>
3	DEBA	Predictions	73	60	50	76	67	59	80	76	72	79	71	67	
		<u>Realizations</u>	<u>73</u>	<u>60</u>	<u>49</u>	<u>77</u>	<u>71</u>	<u>65</u>	<u>80</u>	<u>76</u>	<u>74</u>	79	73	67	<u>70</u>
<u>Equal Weight models</u>															
4	EWc	Predictions	79	68	61	79	73	70	82	78	77	79	69	63	
		<u>Realizations</u>	<u>77</u>	<u>68</u>	<u>62</u>	<u>77</u>	<u>72</u>	<u>68</u>	86	83	81	<u>76</u>	<u>72</u>	<u>65</u>	<u>74</u>
5	EWb	Predictions	71	57	49	75	66	61	79	74	71	77	69	65	
		<u>Realizations</u>	<u>71</u>	<u>58</u>	<u>48</u>	<u>75</u>	<u>67</u>	<u>63</u>	<u>79</u>	<u>75</u>	<u>73</u>	<u>77</u>	<u>69</u>	<u>63</u>	<u>68</u>
<u>Hybrid models</u>															
6	EW/DEBA	Predictions	72	59	50	76	67	62	79	76	73	78	69	67	
		<u>Realizations</u>	<u>73</u>	<u>59</u>	<u>49</u>	<u>77</u>	<u>71</u>	<u>65</u>	<u>79</u>	<u>77</u>	<u>74</u>	79	<u>71</u>	<u>66</u>	<u>70</u>
7	EW/SVb	Predictions	71	58	49	76	67	62	79	76	72	78	68	66	
		<u>Realizations</u>	<u>72</u>	<u>58</u>	<u>49</u>	<u>78</u>	<u>71</u>	<u>65</u>	<u>79</u>	<u>76</u>	<u>74</u>	79	<u>71</u>	<u>66</u>	<u>70</u>
<u>Domran models</u>															
8	DRc	Fit	69	58	51	70	62	58	77	70	66	74	63	57	
		<u>Realizations</u>	<u>69</u>	<u>58</u>	<u>51</u>	<u>69</u>	<u>63</u>	<u>58</u>	<u>77</u>	<u>70</u>	<u>66</u>	<u>73</u>	<u>63</u>	<u>57</u>	<u>65</u>
9	DRb	Fit	70	60	53	74	65	60	78	75	73	75	65	58	
		<u>Realizations</u>	<u>69</u>	<u>60</u>	<u>52</u>	<u>74</u>	<u>66</u>	<u>60</u>	<u>78</u>	<u>75</u>	<u>73</u>	<u>75</u>	<u>65</u>	<u>58</u>	<u>67</u>
<u>Multiple regression</u>															
10	MRc	Predictions	81	70	63	83	79	77	82	78	78	81	73	67	
		<u>Realizations</u>	80	<u>71</u>	<u>65</u>	82	78	75	86	83	81	<u>78</u>	<u>72</u>	<u>66</u>	<u>76</u>
11	MRb	Predictions	64	51	42	64	51	44	72	64	60	66	54	48	
		<u>Realizations</u>	<u>67</u>	<u>54</u>	<u>47</u>	<u>71</u>	<u>62</u>	<u>54</u>	<u>77</u>	<u>73</u>	<u>69</u>	<u>69</u>	<u>66</u>	<u>59</u>	<u>64</u>
		<u>Mean realizations</u>	<u>73</u>	<u>61</u>	<u>53</u>	<u>76</u>	<u>69</u>	<u>63</u>	<u>79</u>	<u>75</u>	<u>72</u>	<u>75</u>	<u>68</u>	<u>62</u>	

Note: **Bold** figures denote largest realization in each column or second largest if MRc is largest.

Figure 1





Figure 2

Percentage correct predictions by different models for conditions specified in Table 5 (cases A, B, C, and D): choosing one of three.








Figure 3

Percentage correct predictions by different models for real data sets specified in Table 12



Appendix A – Correlation between $z_1 = \frac{\varepsilon_b - \varepsilon_a}{\sqrt{2(1 - \rho_{yx}^2)}}$ **and** $z_2 = \frac{\varepsilon_c - \varepsilon_a}{\sqrt{2(1 - \rho_{yx}^2)}}$

Here we prove that $\rho_{z_1, z_2} = 1/2$. First, by definition, and making use of the fact that z_1 and z_2 are standardized normal variables, we have:

$$\rho = \frac{\sigma_{z_1, z_2}}{\sigma_{z_1} \sigma_{z_2}} = \sigma_{z_1, z_2} \quad (\text{A1})$$

$$\sigma \left(\frac{\varepsilon_b - \varepsilon_a}{\sqrt{2(1 - \rho_{yx}^2)}}, \frac{\varepsilon_c - \varepsilon_a}{\sqrt{2(1 - \rho_{yx}^2)}} \right) = \quad (\text{A2})$$

$$\frac{\sigma(\varepsilon_b \varepsilon_c - \varepsilon_b \varepsilon_a - \varepsilon_a \varepsilon_c + \varepsilon_a \varepsilon_a)}{2(1 - \rho_{yx}^2)} \quad (\text{A3})$$

Since ε_a , ε_b , and ε_c are all independent of each other, (A3) can be re-expressed as

$$\rho = \frac{\sigma^2(\varepsilon_a)}{2(1 - \rho_{yx}^2)} = \frac{(1 - \rho_{yx}^2)}{2(1 - \rho_{yx}^2)} = \frac{1}{2} \quad (\text{A4})$$

This result can also be generalized in the following way. First, imagine m different alternatives each of which is characterized by a value of X , i.e., x_l ($l = 1, \dots, m$) and an error term, ε_l ($l = 1, \dots, m$) in similar manner to equations (1) through (3) in the main text. Assume, without loss of generality, that x_l is the largest x value such that we want to calculate the probability that this alternative is better than all the others on the criterion. In this case, all pair-wise correlations between $z_{1l} = \frac{\varepsilon_l - \varepsilon_1}{\sqrt{2(1 - \rho_{yx}^2)}}$ and

$z_{1l+1} = \frac{\varepsilon_{l+1} - \varepsilon_1}{\sqrt{2(1 - \rho_{yx}^2)}}$ for $l = 2, \dots, m$, must – following the logic given above – all equal $1/2$.

Appendix B – Choosing the best of four: the single variable model (SV)

Consider the case of four alternatives, A, B, C, and D with unknown values on the criterion variable, Y . Assume further that A is the preferred choice if $y_a > y_b$, $y_a > y_c$, and $y_a > y_d$. Without loss of generality, let alternative A have the largest value on a single attribute, X , i.e., $x_a > x_b$, $x_a > x_c$, and $x_a > x_d$. Given that it has the largest value, what is the probability that A is the optimal choice, i.e.,

$$P\{(Y_a > Y_b | X_a = x_a > X_b = x_b) \cap (Y_a > Y_c | X_a = x_a > X_c = x_c) \cap (Y_a > Y_d | X_a = x_a > X_d = x_d)\}?$$

Analogically to the case of choice out of three alternatives, and making use of equation (12) in the main text, this probability can be found as

$$P\{(\varepsilon_b - \varepsilon_a < \rho_{yx}(x_a - x_b)) \cap (\varepsilon_c - \varepsilon_a < \rho_{yx}(x_a - x_c)) \cap (\varepsilon_d - \varepsilon_a < \rho_{yx}(x_a - x_d))\} =$$

$$P\left\{\left(z_1 < \frac{\rho_{yx}(x_a - x_b)}{\sqrt{2(1 - \rho_{yx}^2)}}\right) \cap \left(z_2 < \frac{\rho_{yx}(x_a - x_c)}{\sqrt{2(1 - \rho_{yx}^2)}}\right) \cap \left(z_3 < \frac{\rho_{yx}(x_a - x_d)}{\sqrt{2(1 - \rho_{yx}^2)}}\right)\right\} \quad (B1)$$

where z_1 , z_2 , and z_3 are standardized normal variables, jointly distributed with all covariances being equal to $\frac{1}{2}$ (see Appendix A).

Applying the properties of the trivariate normal distribution (i.e., expression (12)), we find the target probability as:

$$\int_{-\infty}^{l_{ab}} \int_{-\infty}^{l_{ac}} \int_{-\infty}^{l_{ad}} \frac{1}{(2\pi)^{3/2} \sqrt{1 - (\rho_{12}^2 + \rho_{13}^2 + \rho_{23}^2) + 2\rho_{12}\rho_{13}\rho_{23}}} e^{-\frac{3}{4}(z_1^2 + z_2^2 + z_3^2) + \frac{1}{2}(z_1 z_2 + z_1 z_3 + z_2 z_3)} dz_1 dz_2 dz_3 =$$

$$\int_{-\infty}^{l_{ab}} \int_{-\infty}^{l_{ac}} \int_{-\infty}^{l_{ad}} \frac{1}{2\pi^{3/2}} e^{-\frac{3}{4}(z_1^2 + z_2^2 + z_3^2) + \frac{1}{2}(z_1 z_2 + z_1 z_3 + z_2 z_3)} dz_1 dz_2 dz_3 \quad (B2)$$

where $l_{ab} = \frac{\rho_{yx}(x_a - x_b)}{\sqrt{2(1 - \rho_{yx}^2)}}; l_{ac} = \frac{\rho_{yx}(x_a - x_c)}{\sqrt{2(1 - \rho_{yx}^2)}}; l_{ad} = \frac{\rho_{yx}(x_a - x_d)}{\sqrt{2(1 - \rho_{yx}^2)}}.$